



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C07D 215/54, A61K 31/47, C07D 211/00, C07D 215/56, C07D 317/00, C07D 401/04, C07D 401/12, C07D 405/12, C07D 413/12, C07D 417/12, C07D 491/10	A1	(11) International Publication Number: WO 00/18740 (43) International Publication Date: 06 April 2000 (06.04.2000)
(21) International Application Number: PCT/US99/22056 (22) International Filing Date: 22 September 1999 (22.09.1999) (30) Priority Data: 09/162,289 29 September 1998 (29.09.1998) US (60) Parent Application or Grant AMERICAN CYANAMID COMPANY [/]; (). WISSNER, Allan [/]; (). TSOU, Hwei-Ru [/]; (). BERGER, Dan, Maarten [/]; (). FLOYD, Middleton, Brawner, Jr. [/]; (). HAMANN, Philip, Ross [/]; (). ZHANG, Nan [/]; (). FROST, Philip [/]; (). MILOWSKY, Arnold, S. ; ().	Published	
(54) Title: SUBSTITUTED 3-CYANOQUINOLINES AS PROTEIN TYROSINE KINASES INHIBITORS (54) Titre: INHIBITEURS DE PROTEINES DE TYPE TYROSINE KINASES A BASE DE 3-CYANOQUINOLINES SUBSTITUEES (57) Abstract <p>This invention provides compounds of formula (1), wherein $R_{\ell 1}$, $G_{\ell 1}$, $G_{\ell 2}$, $R_{\ell 4}$, Z, X and n are defined herein, or a pharmaceutically acceptable salt thereof, which are useful as antineoplastic agents and in the treatment of polycystic kidney disease.</p> (57) Abrégé <p>La présente invention concerne des composés représentés par la formule générale (1) ou certains de leurs sels pharmaceutiquement admis. Dans cette formule, $R_{\ell 1}$, $G_{\ell 1}$, $G_{\ell 2}$, $R_{\ell 4}$, Z, X et n sont tels que décrits dans cette demande internationale. Ces composés, ou les sels considérés, conviennent d'une part comme antinéoplasiques et d'autre part pour le traitement de la polykystose rénale.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 215/54, A61K 31/47, C07D 401/12, 417/12, 215/56, 401/04, 491/10, 405/12, 413/12 // (C07D 491/10, 317:00, 211:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/18740 (43) International Publication Date: 6 April 2000 (06.04.00)</p>
<p>(21) International Application Number: PCT/US99/22056 (22) International Filing Date: 22 September 1999 (22.09.99) (30) Priority Data: 09/162,289 29 September 1998 (29.09.98) US (71) Applicant: AMERICAN CYANAMID COMPANY [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US). (72) Inventors: WISSNER, Allan; 31 Wood Avenue, Ardsley, NY 10502 (US). TSOU, Hwei-Ru; 7 Beverly Place, New City, NY 10956 (US). BERGER, Dan, Maarten; 6 Irion Drive, New City, NY 10956 (US). FLOYD, Middleton, Brawner, Jr.; 5 Babbling Brook Drive, Suffern, NY 10901 (US). HAMANN, Philip, Ross; 9 Alice Street, Gernerville, NY 10923 (US). ZHANG, Nan; 9 New Street, Apartment 2C, Eastchester, NY 10709 (US). FROST, Philip; 4 Emerson Court, Morris Township, NJ 07960 (US). (74) Agents: MILOWSKY, Arnold, S.; American Home Products Corporation, Patent Law Dept.-2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: **SUBSTITUTED 3-CYANOQUINOLINES AS PROTEIN TYROSINE KINASES INHIBITORS**

(57) Abstract

This invention provides compounds of formula (1), wherein R₁, G₁, G₂, R₄, Z, X and n are defined herein, or a pharmaceutically acceptable salt thereof, which are useful as antineoplastic agents and in the treatment of polycystic kidney disease.



(1)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

- 1 -

SUBSTITUTED 3-CYANOQUINOLINES AS PROTEIN TYROSINE KINASES INHIBITORS

BACKGROUND OF THE INVENTION

This invention relates to certain substituted 3-cyano quinoline compounds as well as the pharmaceutically acceptable salts thereof. The compounds of the present invention inhibit the action of certain growth factor receptor protein tyrosine kinases (PTK) and other protein kinases thereby inhibiting the abnormal growth of certain cell types. The compounds of this invention are therefore useful for the treatment of certain diseases that are the result of deregulation of these PTKs. The compounds of this invention are anti-cancer agents and are useful for the treatment of cancer in mammals. In addition, the compounds of this invention are useful for the treatment of polycystic kidney disease in mammals. This invention also relates to the manufacture of said 3-cyano quinolines, their use for the treatment of cancer and polycystic kidney disease, and the pharmaceutical preparations containing them.

Protein tyrosine kinases are a class of enzymes that catalyze the transfer of a phosphate group from ATP to a tyrosine residue located on a protein substrate. Protein tyrosine kinases clearly play a role in normal cell growth. Many of the growth factor receptor proteins function as tyrosine kinases and it is by this process that they effect signaling. The interaction of growth factors with these receptors is a necessary event in normal regulation of cell growth. However, under certain conditions, as a result of either mutation or overexpression, these receptors can become deregulated; the result of which is uncontrolled cell proliferation which can lead to tumor growth and ultimately to the disease known as cancer [Wilks A.F., *Adv. Cancer Res.*, 60, 43 (1993) and Parsons, J.T.; Parsons, S.J., *Important Advances in Oncology*, DeVita V.T. Ed., J.B. Lippincott Co., Phila., 3 (1993)]. Among the growth factor receptor kinases and their proto-oncogenes that have been identified and which are targets of the compounds of this invention are the epidermal growth factor receptor kinase (EGF-R kinase, the protein product of the erbB oncogene), and the product produced by the erbB-2 (also referred to as the neu or HER2) oncogene. Since the phosphorylation event is a necessary signal for cell division to occur and since overexpressed or mutated kinases have been associated with cancer, an inhibitor of

5

- 2 -

10

15

20

25

30

35

40

45

50

55

this event, a protein tyrosine kinase inhibitor, will have therapeutic value for the treatment of cancer and other diseases characterized by uncontrolled or abnormal cell growth. For example, overexpression of the receptor kinase product of the erbB-2 oncogene has been associated with human breast and ovarian cancers [Slamon, D. J., et. al., *Science*, 244, 707 (1989) and *Science*, 235, 1146 (1987)]. Deregulation of EGF-R kinase has been associated with epidermoid tumors [Reiss, M., et. al., *Cancer Res.*, 51, 6254 (1991)], breast tumors [Macias, A., et. al., *Anticancer Res.*, 7, 459 (1987)], and tumors involving other major organs [Gullick, W.J., *Brit. Med. Bull.*, 47, 87 (1991)]. Because of the importance of the role played by deregulated receptor kinases in the pathogenesis of cancer, many recent studies have dealt with the development of specific PTK inhibitors as potential anti-cancer therapeutic agents [some recent reviews: Burke, T.R., *Drugs Future*, 17, 119 (1992) and Chang, C.J.; Geahlen, R.L., *J. Nat. Prod.*, 55, 1529 (1992)]. The compounds of this invention inhibit the kinase activity of EGF-R and are therefore useful for treating certain disease states, such as cancer, that result, at least in part, from deregulation of this receptor. The compounds of this invention are also useful for the treatment and prevention of certain pre-cancerous conditions, such as the growth of colon polyps, that result, at least in part, from deregulation of this receptor.

It is also known that deregulation of EGF receptors is a factor in the growth of epithelial cysts in the disease described as polycystic kidney disease [Du J., Wilson P. D., *Amer. J. Physiol.*, 269(2 Pt 1), 487 (1995); Nauta J., et al., *Pediatric Research*, 37(6), 755 (1995); Gattone V.H., et al., *Developmental Biology*, 169(2), 504 (1995); Wilson P.D., et al., *Eur. J. Cell Biol.*, 61(1), 131, (1993)]. The compounds of this invention, which inhibit the catalytic function of the EGF receptors, are consequently useful for the treatment of this disease.

The mitogen-activated protein kinase (MAPK) pathway is a major pathway in the cellular signal transduction cascade from growth factors to the cell nucleus. The pathway involves kinases at two levels: MAP kinase kinases (MAPKK), and their substrates MAP kinases (MAPK). There are different isoforms in the MAP kinase family. (For review, see Rony Seger and Edwin G. Krebs, *FASEB*, Vol. 9, 726, June 1995). The compounds of this invention can inhibit the action of two of these

5

- 3 -

10

15

20

25

30

35

40

45

50

55

kinases: MEK, a MAP kinase kinase, and its substrate ERK, a MAP kinase. MEK is activated by phosphorylation on two serine residues by upstream kinases such as members of the *raf* family. When activated, MEK catalyzes phosphorylation on a threonine and a tyrosine residue of ERK. The activated ERK then phosphorylates and activates transcription factors in the nucleus, such as *fos* and *jun*, or other cellular targets with PXT/SP sequences. ERK, a p42 MAPK is found to be essential for cell proliferation and differentiation. Over-expression and/or over-activation of Mek or ERK has been found to be associated with various human cancers (For example, Vimala S. Sivaraman, Hsien-yu Wang, Gerard J. Nuovo, and Craig C. Malbon, J. Clin. Invest. Vol. 99, No. 7 April 1997). It has been demonstrated that inhibition of MEK prevents activation of ERK and subsequent activation of ERK substrates in cells, resulting in inhibition of cell growth stimulation and reversal of the phenotype of *ras*-transformed cells (David T. Dudley, Long Pang, Stuart J. Decker, Alexander J. Bridges, and Alan R. Saltiel, PNAS, Vol. 92, 7686, August 1995). Since, as demonstrated below, the compounds of this invention can inhibit the coupled action of MEK and ERK, they are useful for the treatment of diseases such as cancer which are characterized by uncontrolled cell proliferation and which, at least in part, depend on the MAPK pathway.

Epithelial Cell Kinase (ECK) is a receptor protein tyrosine kinase (RPTK) belonging to the EPH (Erythropoietin Producing Hepatoma) family. Although originally identified as an epithelial lineage-specific tyrosine kinase, ECK has subsequently been shown to be expressed on vascular endothelial cells, smooth muscle cells, and fibroblasts. ECK is a type I transmembrane glycoprotein with the extracellular ligand-binding domain consisting of a cysteine-rich region followed by three fibronectin type III repeats. The intracellular domain of ECK possesses a tyrosine kinase catalytic domain that initiates a signal transduction cascade reflecting the ECK function. ECK binds and is subsequently activated by its counter-receptor, Ligand for Eph-Related Kinase (LERK)-1, which is an immediate early response gene product readily inducible in a lineage-unrestricted manner with proinflammatory cytokines such as IL-1 or TNF. Soluble LERK-1 has been shown to stimulate angiogenesis in part by stimulating ECK in a murine model of corneal angiogenesis.

5

- 4 -

10

15

20

25

30

35

40

45

50

55

Unlike their normal counterparts, tumor cells of various lineages constitutively express LERK-1 and this expression can further be upregulated by hypoxia and proinflammatory cytokines. Many of these tumor cells also express ECK at higher levels than their normal counterparts, thereby creating an opportunity for autocrine stimulation via ECK : LERK-1 interaction. The increased expression of both ECK and LERK-1 has been correlated with the transformation of melanomas from the noninvasive horizontal phase of growth into very invasive vertically growing metastatic melanomas. Together, the ECK : LERK-1 interaction is believed to promote tumor growth via its tumor growth promoting and angiogenic effects. Thus, the inhibition of the ECK tyrosine kinase activity mediating signaling cascade induced by its binding and cross-linking to LERK-1 may be therapeutically beneficial in cancer, inflammatory diseases, and hyperproliferative disorders. As is shown below, the compounds of this invention inhibit the tyrosine kinase activity of ECK and are therefore useful for the treatment of the aforementioned disorders.


Growth of most solid tumors is dependent on the angiogenesis involving activation, proliferation and migration of vascular endothelial cells and their subsequent differentiation into capillary tubes. Angiogenization of tumors allows them access to blood-derived oxygen and nutrients, and also provides them adequate perfusion. Hence inhibiting angiogenesis is an important therapeutic strategy in not only cancer but also in a number of chronic diseases such as rheumatoid arthritis, psoriasis, diabetic retinopathy, age-related macular degeneration, and so on. Tumor cells produce a number of angiogenic molecules. Vascular Endothelial Growth Factor (VEGF) is one such angiogenic factor. VEGF, a homodimeric disulfide-linked member of the PDGF family, is an endothelial cell-specific mitogen and is known to cause profound increase in the vascular endothelial permeability in the affected tissues. VEGF is also a senescence-preventing survival factor for endothelial cells. Almost all nucleated tissues in the body possess the capability to express VEGF in response to various stimuli including hypoxia, glucose deprivation, advanced glycation products, inflammatory cytokines, etc. Growth-promoting angiogenic effects of VEGF are mediated predominantly via its signaling receptor Kinase insert Domain containing Receptor (KDR). The expression of KDR is low on most

- 5 -

endothelial cells; however, activation with angiogenic agents results in a significant upregulation of KDR on endothelial cells. Most angiogenized blood vessels express high levels of KDR. KDR is a receptor protein tyrosine kinase with an extracellular VEGF-binding domain consisting of 7 immunoglobulin-like domains and a cytoplasmic domain containing the catalytic tyrosine kinase domain split by a kinase-insert region. Binding to VEGF causes dimerization of KDR resulting in its autophosphorylation and initiation of signaling cascade. Tyrosine kinase activity of KDR is essential for mediation of its functional effects as a receptor for VEGF. Inhibition of KDR-mediated functional effects by inhibiting KDR's catalytic activity is considered to be an important therapeutic strategy in the treatment of angiogenized disease states including cancer. As is shown below, the compounds of this invention inhibit the tyrosine kinase activity of KDR and are therefore useful for the treatment of the aforementioned disease states.

In addition to the above utilities some of the compounds of this invention are useful for the preparation of other compounds of this invention.

The compounds of this invention are certain substituted 3-cyano quinolines. Throughout this patent application, the quinoline ring system will be numbered as indicated in the formula below; the numbering for the quinazoline ring system is also shown :



No 3-cyano quinolines have been reported that have biological activity as inhibitors of protein tyrosine kinases. A 3-cyano quinoline with a 4-(2-methyl anilino) substituent having gastric (H^+/K^+)-ATPase inhibitory activity at high concentrations has been described [Ife R.J., et al., *J. Med. Chem.*, 35(18), 3413 (1992)].

There are quinolines that do not have the 3-cyano substituent and, unlike the compounds of this invention, are unsubstituted at the 4-position but are reported to be inhibitors of protein tyrosine kinases [Gazit A., et al., *J. Med. Chem.*, 39(11), 2170

5

- 6 -

10

15

20

(1996)]. A series of quinolines that have a 3-pyridyl substituent and no substituent at the 4-position have been described as inhibitors of platelet derived growth factor receptor kinase [Dolle R.E., et al., *J. Med. Chem.*, 372, 2627 (1994) and Maguire M.P., et al., *J. Med. Chem.*, 372, 129 (1994)]. The patent applications WO 96/09294 and WO-9813350 describe inhibitors of protein tyrosine kinases that include 4-anilino quinolines with a large variety of substituents on positions 5-8 but which must also have a hydrogen or fluorine atom at position 3. The US patent 5,480,883 describes quinoline derivatives that are inhibitors of protein tyrosine kinases but these derivatives do not have the unique combination of substituents, including the 3-cyano group, contained in the compounds of the present invention. The applications WO-9802434 and WO-9802438 describe quinoline derivatives that are tyrosine kinase inhibitors but these quinolines do not have the important 3-cyano substituent.

25

15

30

35

40

45

50

In addition to quinolines, certain quinazoline derivatives that are similar, in some respects, to the compounds of this invention are known to be inhibitors of protein tyrosine kinases. The application EP-520722 describes 4-anilinoquinazolines that contain simple substituents such as chloro, trifluoromethyl, or nitro groups at positions 5 to 8. The application EP-566226 is similar but with a much larger variety of substituents now allowed at positions 5 to 8. The application WO-9609294 describes compounds with similar substituents at positions 5 to 8 and with the substituent at to 4-position consisting of some polycyclic ring systems. Some simple substituted quinazolines are also described in the applications WO-9524190, WO-9521613, and WO-9515758. The applications EP-602851 and WO-9523141 cover similar quinazoline derivatives where the aryl group attached at position 4 can be a variety of heterocyclic ring structures. The application EP-635498 describes certain quinazoline derivatives that have alkenoylamino and alkynoylamino groups among the substituents at position 6 and a halogen atom at position 7. The application WO-9519774 describes compounds where one or more of the carbon atoms at positions 5-8 can be replaced with heteroatoms resulting in a large variety of bicyclic systems where the left-hand ring is a 5 and 6-membered heterocyclic ring; in addition, a variety of substituents are allowed on the left-hand ring. The application EP-682027-A1 describes certain pyrrolopyrimidine inhibitors of PTKs. The

55


- 7 -

application WO-9519970 describes compounds in which the left-hand aromatic ring of the basic quinazoline structure has been replaced with a wide variety of different heterocyclic rings so that the resulting inhibitors are tricyclic. The application EP-635507 describes quinazolines where an additional 5 or 6-membered heterocyclic ring with optional substitution is fused at positions 5 and 6.

In addition to the aforementioned patent applications, a number of publications describe 4-anilinoquinazolines: Fry, D.W., et. al., *Science*, 265, 1093 (1994), Rewcastle G.W., et. al., *J. Med. Chem.*, 38, 3482 (1995), and Bridges, A.J., et. al., *J. Med. Chem.*, 39, 267, (1996). There are no publications that describe 3-cyano quinolines as PTK inhibitors.

DESCRIPTION OF THE INVENTION

This invention provides a compound of formula 1:



wherein:

X is cycloalkyl of 3 to 7 carbon atoms, which may be optionally substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a pyridinyl, pyrimidinyl, or phenyl ring wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally mono- di-, or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms,

5

- 8 -

10

15

20

25

30

35

40

45

50

55

hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, carboxyalkyl of 2-7 carbon atoms, carboalkoxyalkyl of 3-8 carbon atoms, aminoalkyl of 1-5 carbon atoms, N-alkylaminoalkyl of 2-9 carbon atoms, N,N-dialkylaminoalkyl of 3-10 carbon atoms, N-alkylaminoalkoxy of 2-9 carbon atoms, N,N-dialkylaminoalkoxy of 3-10 carbon atoms, mercapto, and benzoylamino;

Z is -NH-, -O-, -S-, or -NR- ;

R is alkyl of 1-6 carbon atoms, or carboalkyl of 2-7 carbon atoms;


G₁, G₂, R₁, and R₄ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, alkenyloxy of 2-6 carbon atoms, alkynyloxy of 2-6 carbon atoms, hydroxymethyl, halomethyl, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynyloxy of 3-8 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkenoyloxymethyl of 4-9 carbon atoms, alkynyloxymethyl of 4-9 carbon atoms, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulphonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, alkenylsulfonamido of 2-6 carbon atoms, alkynylsulfonamido of 2-6 carbon atoms, hydroxy, trifluoromethyl, trifluoromethoxy, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, amino, hydroxyamino, alkoxyamino of 1-4 carbon atoms, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, phenylamino, benzylamino,

30


5

- 9 -

10




15





with the proviso that either G_1 or G_2 or both G_1 and G_2 must be a radical selected from the group

5

20




25



30

Y is a divalent radical selected from the group consisting of



35

10

R_7 is $-NR_6R_6$, $-J$, $-OR_6$, $-N(R_6)_3^+$, or $-NR_6(OR_6)$;

40

R_7' is $-NR_6(OR_6)$, $-N(R_6)_3^+$, alkenoxy of 1-6 carbon atoms, alkynoxy of 1-6 carbon atoms, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, N-alkyl-N-alkynylamino of 4 to 12 carbon atoms, N-alkenyl-N-alkynylamino of 4 to 12 carbon atoms, or N,N-dialkynylamino of 6-12 carbon atoms with the proviso that the alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

45

M is $>NR_6$, $-O-$, $>N-(C(R_6)_2)_pNR_6R_6$, or $>N-(C(R_6)_2)_p-OR_6$;

W is $>NR_6$, $-O-$ or is a bond;


50

20 Het is a heterocycle selected from the group consisting of morpholine, thiomorpholine, thiomorpholine S-oxide, thiomorpholine S,S-dioxide,

55

- 10 -

piperidine, pyrrolidine, aziridine, pyridine, imidazole, 1,2,3-triazole, 1,2,4-triazole, thiazole, thiazolidine, tetrazole, piperazine, furan, thiophene, tetrahydrothiophene, tetrahydrofuran, dioxane, 1,3-dioxolane ,



tetrahydropyran, and

;

wherein the heterocycle is optionally mono- or di-substituted on carbon or nitrogen with R_6 , optionally mono- or di-substituted on carbon with hydroxy, $-N(R_6)_2$, or $-OR_6$, optionally mono or di-substituted on carbon with the mono-valent radicals $-(C(R_6)_2)_sOR_6$ or $-(C(R_6)_2)_sN(R_6)_2$, or optionally mono or di-substituted on a saturated carbon with divalent radicals $-O-$ or $-O(C(R_6)_2)_sO-$;


R_6 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 1-6 carbon atoms, carboalkyl of 2-7 carbon atoms, carboxyalkyl (2-7 carbon atoms), phenyl, or phenyl optionally substituted with one or more halogen, alkoxy of 1-6 carbon atoms, trifluoromethyl, amino, alkylamino of 1-3 carbon atoms, dialkylamino of 2-6 carbon atoms, nitro, cyano, azido, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, carboxyl, carboalkoxy of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, or alkyl of 1-6 carbon atoms;

R_2 , is selected from the group consisting of


5

- 11 -


10



15




20




25


30



35



40




45


50

55

- 12 -




R₃ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,




with the proviso that at least one of the R₃ groups is selected from the group

- 13 -



R₅ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,



R₈, and R₉ are each, independently, -(C(R₆)₂)_rNR₆R₆, or -(C(R₆)₂)_rOR₆;

J is independently hydrogen, chlorine, fluorine, or bromine;

Q is alkyl of 1-6 carbon atoms or hydrogen;

a = 0 or 1;

g = 1-6;

k = 0-4;

n is 0-1;

p = 2-4;

q=0-4;

r = 1-4;

s = 1-6;

u = 0-4 and v = 0-4, wherein the sum of u+v is 2-4;

or a pharmaceutically acceptable salt thereof,

provided that

- 14 -

when R₆ is alkenyl of 2-7 carbon atoms or alkynyl of 2-7 carbon atoms, such alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

and further provided that

- when Y is -NR₆- and R₇ is -NR₆R₆, -N(R₆)₃, or -NR₆(OR₆), then g = 2-6;
 when M is -O- and R₇ is -OR₆, then p = 1-4;
 when Y is -NR₆-, then k = 2-4;
 when Y is -O- and M or W is -O-, then k = 1-4;
 when W is not a bond with Het bonded through a nitrogen atom, then q = 2-4;
 and when W is a bond with Het bonded through a nitrogen atom and Y is -O- or -NR₆-, then k = 2-4.

The pharmaceutically acceptable salts are those derived from such organic and inorganic acids as: acetic, lactic, citric, tartaric, succinic, maleic, malonic, gluconic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, and similarly known acceptable acids.

The alkyl portion of the alkyl, alkoxy, alkanoyloxy, alkoxymethyl, alkanoyloxymethyl, alkylsulphinyl, alkylsulphonyl, aminoalkyl, N-alkylaminoalkyl, N,N-dialkylaminoalkyl, N-alkylaminoalkoxy, N,N-dialkylaminoalkoxy, alkyl-sulfonamido, carboalkoxy, carboalkyl, carboxyalkyl, carboalkoxyalkyl, alkanoylamino, N-alkylcarbamoyl, and N,N-dialkylcarbamoyl substituents include both straight chain as well as branched carbon chains. The alkenyl portion of the alkenyl, alkenoyloxymethyl, alkenyloxy, alkenylsulfonamido, substituents include both straight chain as well as branched carbon chains and one or more sites of unsaturation and all possible configurational isomers. The alkynyl portion of the alkynyl, alkynoyloxymethyl, alkynylsulfonamido, alkynyloxy, substituents include both straight chain as well as branched carbon chains and one or more sites of unsaturation. Carboxy is defined as a -CO₂H radical. Carboalkoxy of 2-7 carbon atoms is defined as a -CO₂R" radical, where R" is an alkyl radical of 1-6 carbon atoms. Carboxyalkyl is defined as a HO₂C-R'" radical where R'" is a divalent alkyl

5

- 15 -

10

15

20

25

30

35

radical of 1-6 carbon atoms. Carboalkoxyalkyl is defined as a $R''O_2C-R'''$ - radical where R''' is a divalent alkyl radical and where R'' and R''' together have 2-7 carbon atoms. Carboalkyl is defined as a $-COR''$ radical, where R'' is an alkyl radical of 1-6 carbon atoms. Alkanoyloxy is defined as a $-OCOR''$ radical, where R'' is an alkyl radical of 1-6 carbon atoms. Alkanoyloxymethyl is defined as $R''CO_2CH_2$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms. Alkoxymethyl is defined as $R''OCH_2$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms. Alkylsulphinyl is defined as $R''SO$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms. Alkylsulphonyl is defined as $R''SO_2$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms. Alkylsulfonamido, alkenylsulfonamido, alkynylsulfonamido are defined as $R''SO_2NH$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms, an alkenyl radical of 2-6 carbon atoms, or an alkynyl radical of 2-6 carbon atoms, respectively. N-alkylcarbamoyl is defined as $R''NHCO$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms. N,N-dialkylcarbamoyl is defined as $R''R'NCO$ - radical, where R'' is an alkyl radical of 1-6 carbon atoms, R' is an alkyl radical of 1-6 carbon atoms and R' , and R'' may be the same or different. When X is substituted, it is preferred that it is mono-, di-, or tri-substituted, with monosubstituted being most preferred. It is preferred that of the substituents R_1 and R_4 , at least one is hydrogen and it is most preferred that both be hydrogen. It is also preferred that X is a phenyl ring, Z is $-NH-$, and $n = 0$.

40

45

50

Het is a heterocycle, as defined above which may be optionally mono- or di-substituted with R_6 on carbon or nitrogen, optionally mono- or di-substituted on carbon with hydroxy, $-N(R_6)_2$, or $-OR_6$, optionally mono or di-substituted on carbon with $-(C(R_6)_2)_sOR_6$ or $-(C(R_6)_2)_sN(R_6)_2$, and optionally mono or di-substituted on a saturated carbon with divalent radicals $-O-$ or $-O(C(R_6)_2)_sO-$ (carbonyl and ketal groups, respectively); in some cases when Het is substituted with $-O-$ (carbonyl), the carbonyl group can be hydrated. Het may be bonded to W when $q = 0$ via a carbon atom on the heterocyclic ring, or when Het is a nitrogen containing heterocycle which also contains a saturated carbon-nitrogen bond, such heterocycle may be bonded to carbon, via the nitrogen when W is a bond. When $q = 0$ and Het is

55


- 16 -

a nitrogen containing heterocycle which also contains an unsaturated carbon-nitrogen bond, that nitrogen atom of the heterocycle may be bonded to carbon when W is a bond and the resulting heterocycle will bear a positive charge. When Het is substituted with R₆, such substitution may be on a ring carbon, or in the case of a nitrogen containing heterocycle, which also contains a saturated carbon-nitrogen, such nitrogen may be substituted with R₆ or in the case of a nitrogen containing heterocycle, which also contains an unsaturated carbon-nitrogen, such nitrogen may be substituted with R₆ in which case the heterocycle will bear a positive charge. Preferred heterocycles include pyridine, 2,6-disubstituted morpholine, 2,5-disubstituted thiomorpholine, 2-substituted imidazole, substituted thiazole, thiazolidine, N-substituted imidazole, N-substituted 1,4-piperazine, N-substituted piperidine, dioxane, 1,3-dioxolane, and N-substituted pyrrolidine.

The compounds of this invention may contain one or more asymmetric carbon atoms; in such cases, the compounds of this invention include the individual diastereomers, the racemates, and the individual R and S enantiomers thereof. Some of the compounds of this invention may contain one or more double bonds; in such cases, the compounds of this invention include each of the possible configurational isomers as well as mixtures of these isomers.

The compounds having formula 1 and their salts may be prepared by a process which comprises

(a) reacting a compound having the formula



where R₁, G₁, G₂, R₄, Z, n and X are as defined above with a dehydrating agent so as to convert the aminocarbonyl group into a cyano group, or

- 17 -

(b) reacting a compound having the formula



or a salt thereof with a compound having the formula



where Q is a leaving group and A_1 , A_2 and A_3 are such that $A_1-NH-A_2-A_3$ is a compound conforming with formula 1; or

(c) reacting a compound having the formula



or a salt thereof with a compound having the formula



where Q is as defined above and A_4 and A_5 are such that A_4-O-A_5 is a compound conforming with formula 1; or

(d) adding an acid to a compound having formula 1 so that an acid addition salt is prepared.

The preparation of the compounds and intermediates of this invention encompassed by Formula 5 is described below in Flowsheet 1 where Z and n are as described above and X' is cycloalkyl of 3 to 7 carbon atoms, which may be optionally substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a pyridinyl, pyrimidinyl, or phenyl ring wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally mono- di-, or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms,

5

- 18 -

10

alkynoylamino of 3-8 carbon atoms, carboxyalkyl of 2-7 carbon atoms, aminomethyl, N-alkylaminomethyl of 2-7 carbon atoms, N,N-dialkylaminomethyl of 3-7 carbon atoms, mercapto, and benzoylamino;

15

R₁', R₂', R₃', and R₄' are each, independently, hydrogen, halogeno, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, alkenyloxy of 2-6 carbon atoms, alkynyloxy of 2-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulphonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, trifluoromethyl, cyano, nitro, carboxy, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, alkoxyamino of 1-4 carbon atoms, dialkylamino of 2 to 12 carbon atom, N,N-dialkylaminoalkyl of 3-14 carbon atoms, phenylamino, benzylamino, N-alkylcarbamoyl of 1-6 carbon atoms, N,N-dialkylcarbamoyl of 2-12 carbon atoms. According to the sequence of reaction outlined in flowsheet 1, a quinoline-3-carboxylic acid ester of Formula 2 is

25

30

hydrolyzed with base to furnish a carboxylic acid of Formula 3. The carboxylic acid group of 3 is converted to an acyl imidazole by heating it with carbonyldiimidazole in an inert solvent such as dimethylformamide (DMF) followed by the addition of ammonia to give the amide 4. Dehydration of the amide functional group with a dehydrating agent such as trifluoroacetic anhydride in pyridine, phosphorous pentoxide in an inert solvent, or the like gives the 3-cyano quinolines, 5, of this invention. In those cases where any of the intermediates have an asymmetric carbon atom, they can be used as the racemate or as the individual R or S enantiomers in which case the compounds of this invention will be in the racemic or R and S optically active forms, respectively. The quinoline-3-carboxylic acid esters of Formula 2, the quinoline-3-carboxylic acids of Formula 3, and the quinoline-3-carboxylic amides of Formula 4 needed to prepare the compounds of this invention are either already known to the art or can be prepared by procedures known in the art as detailed in the following references:

35

40

45

50


Sarges, Reinhard; Gallagher, Andrea; Chambers, Timothy J.; Yeh, Li An, *J. Med. Chem.*, 36, 2828 (1993); Savini, Luisa; Massarelli, Paola; Pellerano, Cesare; Bruni, Giancarlo, *Farmaco*, 48(6), 805 (1993); Ife, Robert J.; Brown, Thomas H.; Keeling,

55

- 19 -

David J.; Leach, Colin, *J. Med. Chem.*, 35, 3413 (1992); Hanifin, J. William;
 Capuzzi, Rosemary; Cohen, Elliott, *J. Med. Chem.*, 12(5), 1096 (1969); Marecki,
 Paul E.; Bambury, Ronald E., *J. Pharm. Sci.*, 73(8), 1141 (1984); Pellerano, C.;
 Savini, L.; Massarelli, P.; Bruni, G.; Fiaschi, A. I., *Farmaco*, 45(3), 269, (1990);
 Marecki, Paul E.; Bambury, Ronald E., *J. Pharm. Sci.*, 73(8), 114 (1984); patent
 application WO 8908105; US patent 4343804; US patent 3470186.

FLOWSHEET 1



10

The preparation of the compounds of this invention encompassed by Formula 12 is
 described below in Flowsheet 2 where X, Z, n, R₁, G₂, G₁, and R₃ are as described

55

5

- 20 -

10

15

20

25

30

35

40

45

50


55

above. The substituted aniline of Formula 6 is heated with or without a solvent with the reagent 7 to give intermediate 8 as a mixture of isomers. Thermolysis of 8 in a high boiling solvent such as diphenyl ether at 200-350°C gives the 3-cyano quinolones of Formula 9; these intermediates may also exist in the 4-hydroxy quinoline tautomeric form. In those cases where R_4 is a hydrogen atom, the intermediates 9 may be formed as a mixture of two regioisomers. These isomers can be separated by methods well known in the art including, but not limited to, fractional crystallization and chromatographic methods. The separated isomers can then be converted separately to the compounds of the invention. Alternatively, the isomers can be separated at a later stage of the synthesis. Heating compounds 9 with or without solvent with a chlorinating agent such as phosphorous oxychloride or phosphorous pentachloride gives the 4-chloro-3-cyano quinolines of Formula 10. Condensation of 10 with a nucleophilic amine, aniline, mercaptan, thiophenol, phenol, or alcohol reagent of Formula 11 gives the 3-cyano quinolines intermediates of Formula 12; this condensation can be accelerated by heating the reaction mixture or by using basic catalysts such as trialkylamines, sodium hydride in an inert solvent, sodium or potassium alkoxides in an alcohol solvents, and the like. In those cases where the substituents may contribute an asymmetric carbon atom, the intermediates can be used as the racemate or as the individual R or S enantiomers in which case the compounds of this invention will be in the racemic or R and S optically active forms, respectively. In cases where the substituents may contribute more than one asymmetric carbon atoms, diastereomers may be present; these can be separated by methods well known in the art including, but not limited to, fractional crystallization and chromatographic methods. In those cases where R_1 , G_2 , G_1 , and R_4 moieties contain primary or secondary amino groups, the amino groups may first have to be used in protected form prior to reaction with reagent 7. Suitable protecting groups include, but are not limited to, tert-butoxycarbonyl (BOC) and benzyloxycarbonyl (CBZ) protecting groups. The former protecting group can be removed from the final products of Formula 12 by treatment with an acid such as trifluoroacetic acid while the latter protecting group can be removed by catalytic hydrogenation. In those cases where the R_1 , G_2 , G_1 , and R_4 moieties contain hydroxyl groups, the hydroxyl groups

- 21 -

may first have to be used in protected form prior to reaction with reagent 7. Suitable protecting groups include, but are not limited to, t-butyldimethylsilyl, tetrahydropyranyl, or benzyl protecting groups. The first two protecting groups can be removed from the final products of formula 12 by treatment with an acid such as acetic acid or hydrochloric acid while the latter protecting group can be removed by catalytic hydrogenation.

FLOWSHEET 2



5

- 22 -

10

15

20

25

30

35

40

45


50

55





The preparation of intermediate **15** (identical to intermediate **9** of Flowsheet 2) can also be prepared as describe below in Flowsheet 3. Heating the substituted aniline of Formula **13** with dimethylformamide dimethyl acetal with or without a solvent gives intermediates for Formula **14**. The reaction of **14** with the lithium anion of acetonitrile prepared using a base such as n-butyl lithium or the like in an inert solvent gives the 3-cyano quinolones, **15**, or the 3-cyano-4-hydroxy quinoline tautomers thereof which can be converted to the compounds of this invention. In those cases where R_1 , G_2 , G_1 , and R_4 moieties contain primary or secondary amino groups, the amino groups may first have to be used in protected form. Suitable protecting groups include, but are not limited to, tert-butoxycarbonyl (BOC) and benzyloxycarbonyl (CBZ) protecting groups. The former protecting group can be removed from the final products of Formula **15** by treatment with an acid such as trifluoroacetic acid while the latter protecting group can be removed by catalytic hydrogenation. In those cases where the R_1 , G_2 , G_1 , and R_4 moieties contain hydroxyl groups, the hydroxyl groups may first have to be used in protected form. Suitable protecting groups include, but are not limited to, t-butyldimethylsilyl, tetrahydropyranyl, or benzyl protecting groups. The first two protecting groups can be removed from the final products of formula **15** by treatment with an acid such as acetic acid or hydrochloric acid while the latter protecting group can be removed by catalytic hydrogenation.

- 23 -

FLOWSHEET 3



- 5 The preparation of the compounds of this invention encompassed by Formula 24 is described below in Flowsheet 4 wherein R_1 , G_2 , R_4 , Z , n , and X are defined. R_{10} is alkyl of 1-6 carbon atoms (preferably isobutyl). R_2' is a radical selected from the group consisting of:



wherein R₆, R₃, R₅, J, s, r, u, and v are defined. According to the reactions outlined in Flowsheet 4, a 4-chloro-3-cyano-6-nitroquinoline, 16, can be reacted with an amine or aniline 17 by heating in an inert solvent such as tetrahydrofuran, butanol, or methoxyethanol to give compounds of Formula 20 where Z is -NH-. The reaction of 16 with a mercaptan or thiophenol 18 in an inert solvent can be accomplished using a base such as sodium hydride to give compounds of Formula 20 where Z is -S-. The reaction of 16 with an alcohol or phenol 19 in an inert solvent can be accomplished using a base such as sodium hydride to give compounds of Formula 20 where Z is -O-. Compounds of Formula 20 can be reduced to a 6-amino-3-cyano-quinoline, 21, using a reducing agent such as sodium hydrosulfite in a two phase system consisting

5

- 25 -

10

15

20

25

30

35

40

45


50

55


of tetrahydrofuran and water in the presence of a small amount of phase transfer catalyst or by using iron in refluxing protic solvents containing acetic acid or ammonium chloride. Acylation of **21** with either an acid chloride of Formula **22** or a mixed anhydride of Formula **23** (which is prepared from the corresponding carboxylic acid) in an inert solvent such as tetrahydrofuran (THF) in the presence of an organic base such as pyridine, triethylamine, diisopropylethylamine, or N-methyl morpholine gives the compounds of this invention of Formula **24**. In those cases where **22** or **23** have an asymmetric carbon atom, they can be used as the racemate or as the individual R or S enantiomers in which case the compounds of this invention will be in the racemic or R and S optically active forms, respectively. In those cases, where the R₂' contains primary or secondary amino groups, the amino groups will first have to be protected prior to anhydride or acid chloride formation. Suitable protecting groups include, but are not limited to, tert-butoxycarbonyl (BOC) and benzyloxycarbonyl (CBZ) protecting groups. The former protecting group can be removed from the final products of Formula **24** by treatment with an acid such as trifluoroacetic acid while the latter protecting group can be removed by catalytic hydrogenation. In those cases where the R₂' contains hydroxyl groups, the hydroxyl groups may first have to be protected prior to anhydride or acid chloride formation. Suitable protecting groups include, but are not limited to, t-butyldimethylsilyl, tetrahydropyranyl, or benzyl protecting groups. The first two protecting groups can be removed from the final products of Formula **24** by treatment with an acid such as acetic acid or hydrochloric acid while the latter protecting group can be removed by catalytic hydrogenation. In those cases, in intermediates **17**, **18**, or **19** where X contains primary or secondary amino groups or hydroxyl groups, it may be necessary to protect these groups prior to the reaction with **16**. The same amine or alcohol protecting groups describe above can be used and they can be removed from the products **24** as previously described.

- 26 -

FLOWSHEET 4



By using methods similar to that describe above in Flowsheet 4, the intermediates 25 can be converted to the compounds of this invention, 26.



5

- 27 -

10

15

20

In order to prepare the compounds of this invention, certain amines are required. Some representative amines are shown below in List A wherein R₆, p, and r are as defined above. These amines are available commercially, are known in the chemical literature, or can be prepared by simple procedures that are well known in the art. In some cases, these amines may have an asymmetric carbon atoms; they can be used as the racemate or they can be resolved and used as the individual R or S enantiomers in which case the compounds of this invention will be in the racemic or optically active forms, respectively. Throughout this application in the Flowsheets shown below, these amines, and other similar amines, will be represented by the generic structure of the formula:

25

$(R')_2NH$, wherein this formula can represent a primary or secondary amine.

30

35


40

45

50

55

List A




- 29 -

In order to prepare the compounds of this invention certain alcohols are required. Some representative alcohols are shown below in List B wherein R_6 , p , and r are as defined above. These alcohols are available commercially, are known in the chemical literature, or can be prepared by simple procedures that are well known in the art. In some cases, these alcohols may have an asymmetric carbon atoms; they can be used as the racemate or they can be resolved and used as the individual R or S enantiomers in which case the compounds of this invention will be in the racemic or optically active forms, respectively. Throughout this application in the Flowsheets shown below, these alcohols, and other similar alcohols, will be represented by the generic structure of the formula:



List B



5

- 30 -

10

15

20

25

30

35

40

45

50

55

In order to prepare some of the compounds of this invention certain mixed anhydrides of Formulas 31, 34, and 38 are required; these are prepared as outlined below in Flowsheet 5-6 wherein R₆, R₁₀, X, Z, n, and s are as defined above. J' is a halogen atom chlorine, bromine, or iodine, or is a tosylate (p-toluenesulfonate) or mesylate (methanesulfonate) group. The reaction of 27 with an amine of List A is accomplished by heating in an inert solvent such as tetrahydrofuran or N,N-dimethylformamide, or using potassium or cesium carbonate in acetone. The temperature and duration of the heating will depend on the reactivity of 27; longer reaction times and higher temperatures may be required when s is greater than 1.





10 Treatment of 28 with an alkyl lithium reagent followed by quenching with an atmosphere of dry carbon dioxide furnishes the carboxylic acids of formula 29. These can be converted to mixed anhydrides of Formula 31 using a reagent such as isobutylchloroformate in an inert solvent such as tetrahydrofuran in the presence of a base such as N-methylmorpholine. These anhydrides can then be used to prepare the compounds of this invention as described above in Flowsheet 4. The reaction of 27

15 with an alcohol of List B is accomplished using sodium hydride or other non-nucleophilic base such as potassium or cesium carbonate in an inert solvent such as tetrahydrofuran, acetone, or N,N-dimethylformamide. In some cases, the alcohol of List B can also be the solvent of the reaction. Treatment of 32 with an alkyl lithium

20 reagent followed by quenching with an atmosphere of dry carbon dioxide furnishes the carboxylic acids of formula 33. These can be converted to mixed anhydrides Formula 34 using a reagent such as isobutylchloroformate in an inert solvent such as tetrahydrofuran in the presence of a base such as N-methylmorpholine. These anhydrides can then be used to prepare the compounds of this invention as described

25 above in Flowsheet 4.

FLOWSHEET 5




As outline in Flowsheet 6 below wherein R_1 , G_2 , R_4 , R_6 , R_{10} , X , Z , n , and s are as defined above, alcohols 35 can be protected with a *t*-butyl dimethylsilyl protecting group by the reaction with the respective silyl chloride in methylene chloride in the presence of triethylamine and 4-*N,N*-dimethylamino pyridine (DMAP). The resulting protected alcohols, 36, are converted to the acetylenic

- 32 -

Grignard reagents which, in turn, are maintained under an atmosphere of dry carbon dioxide to give the carboxylic acids **37**. As described above these are converted to the mixed anhydrides **38** which on reaction with the 6-amino-3-cyanoquinoline **39** gives **40**. In the final step of the sequence, the silyl protecting group is removed by treating with acid in a protic solvent mixture to give the compounds represented by Formula **41**.

FLOWSHEET 6



5

- 33 -

10

15

20

25

30

35

40

45


50

55

Compounds of this invention are also prepared as shown below in Flowsheet 7 wherein R₁, G₂, R₄, R₆, R₁₀, X, Z, n, and s are as defined above. J' is a halogen atom chlorine, bromine, or iodine, or is a tosylate or mesylate group. Treatment of **42** with an alkyl lithium reagent at low temperature followed by quenching with an atmosphere of dry carbon dioxide furnishes the carboxylic acids of formula **43**. These can be converted to mixed anhydrides of Formula **44** using a reagent such as isobutylchloroformate in an inert solvent such as tetrahydrofuran in the presence of a base such as N-methylmorpholine. These anhydrides can then be used to prepare the compounds of this invention as by the reaction with the 6-amino-3-cyanoquinolines **45** described above in the Flowsheets. The reaction of **46** with an alcohol of List **B** is accomplished using sodium hydride or other non-nucleophilic base in an inert solvent such as tetrahydrofuran or N,N-dimethylformamide to give the compounds of this invention represented by **47**. In some cases, the alcohol of List **B** can also be the solvent of the reaction. The reaction of **46** with an amine of List **A** gives the compounds of this invention represented by **48** is accomplished by heating in an inert solvent such as tetrahydrofuran or N,N-dimethylformamide, or using potassium or cesium carbonate in acetone. The temperature and duration of the heating will depend on the reactivity of **46**; longer reaction times and higher temperatures may be required when s is greater than 1.


- 34 -

FLOWSHEET 7



Using methods similar to that summarized above, **45b** can be converted to **47b** or **48b**.

- 35 -



Other carboxylic acid chlorides and anhydrides needed to prepare some of the compounds of this invention are prepared as shown below in Flowsheet 8 wherein

5 R₆, R₃, R₁₀, X, Z, J', n, and s are as defined above. Q' is an alkyl group of 1-6 carbon atoms. The esters **49**, **53**, or **57** can be hydrolyzed with a base such as barium hydroxide to give the respective carboxylic acid **50**, **54**, or **58**. These acid can be


10 converted to the respective carboxylic acid chlorides **51** or **56** by using oxalyl chloride and catalytic N,N-dimethylformamide in an inert solvent or respective mixed anhydrides **55** or **59** by using isobutyl chloroformate and an organic base such as N-methylmorpholine. The leaving group in compounds represented by Formula **52** can

15 be displaced by the amines of List A or the alcohols of List B by using procedures previously described to give the intermediates **57** and **53**, respectively. These carboxylic acid chlorides **51** and **56** and these anhydrides **55** and **59** can be used to

20 prepare some of the compounds of this invention by using the methods outlined herein above in the Flowsheets.


- 36 -

FLOWSHEET 8




- 37 -

By using the methods identical to those outlined above in Flowsheet 8, it is possible to prepare the analogous carboxylic acid chlorides and anhydrides given below in List C wherein R₆, R₃, p, and s are as previously defined. G is the radical:




and A is the radical:



wherein $-\text{N}(\text{R}')_2$ is derived from the amines of List A, $-\text{OR}'$ are derived from the alcohols of List B, and J' is a leaving group as defined previously. By making use of these carboxylic acid chlorides and anhydrides, by following the methods summarized in the above in Flowsheets, and by pursuing the details described in the examples given below, many of the compounds of this invention can be prepared.

- 38 -

LIST C




Compounds of this invention represented by Formulas **62-63** can be prepared as shown in Flowsheet **9** wherein R₁, G₂, R₄, R₆, R₃, R₁₀, X, Z, J', n, and s are as defined above. The reaction of the carboxylic acid chlorides **60** and the 6-amino-3-cyanoquinolines **61** using an organic base in an inert solvent gives the compounds of this invention represented by Formula **62**. The reaction of **62** with an alcohol of List **B** is accomplished using sodium hydride or other non-nucleophilic base such as potassium or cesium carbonate in an inert solvent such as tetrahydrofuran, acetone, or N,N-dimethylformamide to give the compounds of this invention represented by **63**.

- 39 -


In some cases, the alcohol of List B can also be the solvent of the reaction. The reaction of **62** with an amine of List A to give the compounds of this invention represented by **64** is accomplished by heating in an inert solvent such as tetrahydrofuran or N,N-dimethylformamide. The temperature and duration of the heating will depend on the reactivity of **62**; longer reaction times and higher temperatures may be required when s is greater than 1. In addition, by using this method, the carboxylic acid chlorides and mixed anhydrides listed in List C can be used to prepare the analogous compounds of this invention.

FLOWSHEET 9




- 40 -

By applying the methods summarized above, **61b** can be converted to **63b** and **64b** via the intermediate **62b**.



- 41 -



The reaction of **62** or **62b** with a nitrogen containing heterocycle HET which also contains an unsaturated carbon-nitrogen bond is accomplished by refluxing in an inert solvent and gives the compounds of this invention **64c** and **64d**, respectively where the compound bears a positive charge. The counter anion J⁻ can be replaced with any other pharmaceutically acceptable anion using the appropriate ion exchange resin.



Some of the compounds of this invention can be prepared as outline below in Flowsheet 10 wherein R₁, G₂, R₃, R₄, R₆, R₁₀, X, Z, J⁻, n, and r are as defined above. The acetylenic alcohols **65** can be coupled to the halides, mesylates, or tosylates **66** using a base such as sodium hydride in an inert solvent such as tetrahydrofuran. The resulting acetylene, **67**, is then treated with an alkyl lithium reagent at low temperature. Maintaining the reaction under an atmosphere of carbon dioxide then gives the carboxylic acids **68**. These, in turn, are reacted with the 6-amino-3-cyanoquinolines, **69**, via the mixed anhydrides to give the compounds of this invention represented by Formula **70**. Alternatively, the intermediates **67** can be prepared starting with an alcohol **71** by first treating it with a base such as sodium hydride in an inert solvent such as tetrahydrofuran and then adding an acetylene **72** that has an appropriate leaving group. In a similar manner, the amino alcohols






- 42 -

represented by the formula: $(R_6)_2N-(C(R_6)_2)_r-OH$ by reacting with 72, and applying the chemistry of Flowsheet 10, can be converted to the compounds of this invention represented by the formulas:




- 43 -

FLOWSHEET 10




- 44 -

By applying similar methods as described above, **69b** can be converted to the compounds of this invention represented by **70b**.




The compounds of this invention represented by Formula **76** and **77** are prepared as shown below in Flowsheet **11** wherein R_1 , R_3 , R_4 , R_6 , and n defined above and the amines $\text{HN}(\text{R}'')_2$ are selected from the group:



Refluxing **73** and **74** in an a solvent such as ethanol gives the intermediate **75** which can react with an amine in refluxing ethanol to give the compounds of this invention represented by Formula **76**. Treating **75** with an excess of a sodium alkoxide in an inert solvent or a solvent from which the alkoxide is derived gives the compounds of this invention of Formula **77**.



- 45 -

FLOWSHEET 11



In a manner similar to that described above, 74b can be converted to 76b or 77b.


- 46 -

**74b****76b****77b**


Compounds of this invention represented by Formula **83** can be prepared as shown in Flowsheet **12** wherein R_1 , G_2 , R_4 , R_6 , R_3 , R_{10} , X , Z , n , and r are as defined above. The reaction of the mercapto carboxylic acids **78** with the reagents **79** give the compounds represented by Formula **80**. Alternatively, **80** can be prepared from the mercaptan R_3SH using the mercapto acid **78**, triethylamine and 2,2'-dipyridyl disulfide. Mixed anhydride formation to give **81** followed by condensation with the 6-amino-3-cyanoquinolines **82** give the compounds of this invention.

10

FLOWSHEET 12



5 By applying similar methods as described above **82b** can be converted to **83b**.



5

- 48 -

10

15

20

Compounds of this invention represented by Formulas **86-88** can be prepared as shown in Flowsheet **13** wherein R_1 , G_2 , R_1 , R_4 , R_5 , J , X , Z , and n are as defined above. Q' is alkyl of 1-6 hydrogen atoms, alkoxy of 1-6 hydrogen atoms, hydroxy, or hydrogen. Akylation of **84** with the 6-amino-3-cyanoquinolines **85** can be accomplished by heating in an inert solvent such as N,N-dimethylformamide using a base such as potassium carbonate to give the compounds of this invention represented by the Formula **86**. When Q' is alkoxy, the ester group can be hydrolyzed to an acid using a base such as sodium hydroxide in methanol. In a similar manner, by using intermediates **89** and **90**, the compounds of this invention represented by Formulas **87** and **88** can be prepared, respectively.

25

30

35

40







45

50

55





- 49 -

FLOWSHEET 13



- 50 -


By applying similar methods as described above **85b** can be converted to **86b-88b**.



Compounds of this invention represented by Formula **93** can be prepared as shown in Flowsheet **14** wherein R_1 , G_2 , R_1 , R_4 , R_5 , X , Z , and n are as defined above. The reaction of reagent **91** with the 6-amino-3-cyanoquinolines **92** is accomplished using an excess of an organic base such as triethylamine and an inert solvent such as tetrahydrofuran to give compounds of this invention represented by Formula **93**.

- 51 -


FLOWSHEET 14



Compounds of this invention represented by Formula 96 can be prepared as shown in Flowsheet 15 wherein R_1 , G_1 , R_1 , R_4 , R_5 , R_6 , W , Het , X , Z , k , and n are as defined above by the Mitsunobu reaction of phenol 94 and an alcohol 95 in an inert solvent. Alternatively, the Mitsunobu reaction can be applied to compound 97 to give 98. This compound can be converted to 96 as described above in Flowsheet 4. The heterocycle can be introduced at the 6-position by using the corresponding compounds where G_1 is hydroxy and G_2 is located at the 7-position.

- 52 -

FLOWSHEET 15



There are certain functional group manipulations that are useful to prepare the compounds of this invention that can be applied to various intermediate 3-cyanoquinolines as well as to the final compounds of this invention. These manipulations refer to the substituents R_1 , G_1 , G_2 , or R_4 that are located on the 3-cyanoquinolines shown in the above Flowsheets. Some of these functional group manipulations are described below:

Where one or more of R_1 , G_1 , G_2 , or R_4 is a nitro group, it can be converted to the corresponding amino group by reduction using a reducing agent such as iron in acetic acid or by catalytic hydrogenation. Where one or more of R_1 , G_1 , G_2 , or R_4 is an amino group, it can be converted to the corresponding dialkylamino group of 2 to 12 carbon atoms by alkylation with at least two equivalents of an alkyl halide of 1 to

5

- 53 -

10

15

20

25

30

35

40

45

50

55

6 carbon atoms by heating in an inert solvent or by reductive alkylation using an aldehyde of 1 to 6 carbon atoms and a reducing agent such as sodium cyanoborohydride. Where one or more of R₁, G₁, G₂, or R₄ is a methoxy group, it can be converted to the corresponding hydroxy group by reaction with a demethylating agent such as boron tribromide in an inert solvent or by heating with pyridinium chloride with or without solvent. Where one or more of R₁, G₁, G₂, or R₄ is an amino group, it can be converted to the corresponding alkylsulfonamido, alkenylsulfonamido, or alkynylsulfonamido group of 2 to 6 carbon atoms by the reaction with an alkylsulfonyl chloride, alkenylsulfonyl chloride, or alkynylsulfonyl chloride, respectively, in an inert solvent using a basic catalyst such as triethylamine or pyridine. Where one or more of R₁, G₁, G₂, or R₄ is an amino group, it can be converted to the corresponding alkylamino group of 1 to 6 carbon atoms by alkylation with one equivalent of an alkyl halide of 1 to 6 carbon atoms by heating in an inert solvent or by reductive alkylation using an aldehyde of 1 to 6 carbon atoms and a reducing agent such as sodium cyanoborohydride in a protic solvent such as water or alcohol, or mixtures thereof. Where one or more of R₁, G₁, G₂, or R₄ is hydroxy, it can be converted to the corresponding alkanoyloxy group of 1-6 carbon atoms by reaction with an appropriate carboxylic acid chloride, anhydride, or mixed anhydride in an inert solvent using pyridine or a trialkylamine as a catalyst. Where one or more of R₁, G₁, G₂, or R₄ is hydroxy, it can be converted to the corresponding alkenoyloxy group of 1-6 carbon atoms by reaction with an appropriate carboxylic acid chloride, anhydride, or mixed anhydride in an inert solvent using pyridine or a trialkylamine as a catalyst. Where one or more of R₁, G₁, G₂, or R₄ is hydroxy, it can be converted to the corresponding alkynoyloxy group of 1-6 carbon atoms by reaction with an appropriate carboxylic acid chloride, anhydride, or mixed anhydride in an inert solvent using pyridine or a trialkylamine as a catalyst. Where one or more of R₁, G₁, G₂, or R₄ is carboxy or a carboalkoxy group of 2-7 carbon atoms, it can be converted to the corresponding hydroxymethyl group by reduction with an appropriate reducing agent such as borane, lithium borohydride, or lithium aluminum hydride in an inert solvent; the hydroxymethyl group, in turn, can be converted to the corresponding halomethyl

5

- 54 -

10

15

20

25

30

35

40

45

50

55

group by reaction in an inert solvent with a halogenating reagent such as phosphorous tribromide to give a bromomethyl group, or phosphorous pentachloride to give a chloromethyl group. The hydroxymethyl group can be acylated with an appropriate acid chloride, anhydride, or mixed anhydride in an inert solvent using pyridine or a trialkylamine as a catalyst to give the compounds of this invention with the corresponding alkanoyloxymethyl group of 2-7 carbon atoms, alkenoyloxymethyl group of 2-7 carbon atoms, or alkynoyloxymethyl group of 2-7 carbon atoms. Where one or more of R₁, G₁, G₂, or R₄ is a halomethyl group, it can be converted to an alkoxymethyl group of 2-7 carbon atoms by displacing the halogen atom with a sodium alkoxide in an inert solvent. Where one or more of R₁, G₁, G₂, or R₄ is a halomethyl group, it can be converted to an aminomethyl group, N-alkylaminomethyl group of 2-7 carbon atoms or N,N-dialkylaminomethyl group of 3-14 carbon atoms by displacing the halogen atom with ammonia, a primary, or secondary amine, respectively, in an inert solvent.

15 In addition to the methods described herein above, there a number of patent applications that describe methods that are useful for the preparation of the compounds of this invention. Although these methods describe the preparation of certain quinazolines, they are also applicable to the preparation of correspondingly substituted 3-cyanoquinolines. The chemical procedures described in the application
20 WO-9633981 can be used to prepare the 3-cyanoquinoline intermediates used in this invention wherein R₁, G₁, G₂, or R₄ are alkoxyalkylamino groups. The chemical procedures described in the application WO-9633980 can be used to prepare the 3-cyanoquinoline intermediates used in this invention wherein R₁, G₁, G₂, or R₄ are aminoalkylalkoxy groups. The chemical procedures described in the application WO-
25 9633979 can be used to prepare the 3-cyanoquinoline intermediates used in this invention wherein R₁, G₁, G₂, or R₄ are alkoxyalkylamino groups. The chemical procedures described in the application WO-9633978 can be used to prepare the 3-cyanoquinoline intermediates used in this invention wherein R₁, G₁, G₂, or R₄ are aminoalkylamino groups. The chemical procedures described in the application WO-
30 9633977 can be used to prepare the 3-cyanoquinoline intermediates used in this

5

- 55 -

10

invention wherein R₁, G₁, G₂, or R₄ are aminoalkylalkoxy groups. Although the above patent applications describe compounds where the indicated functional group have been introduced onto the 6-position of a quinazoline, the same chemistry can be used to introduce the same groups unto positions occupied by the R₁, G₁, G₂, and R₄

15

5 substituents of the compounds of this invention

20

Representative compounds of this invention were evaluated in several standard pharmacological test procedures that showed that the compounds of this invention possess significant activity as inhibitors of protein tyrosine kinases, and are antiproliferative agents. Based on the activity shown in the standard pharmacological

10 test procedures, the compounds of this invention are therefore useful as antineoplastic agents. The test procedures used and results obtained are shown below.

25

Inhibition of Epidermal Growth Factor Receptor Kinase (EGF-R)

15

using recombinant enzyme

30

Representative test compounds were evaluated for their ability to inhibit the phosphorylation of the tyrosine residue of a peptide substrate catalyzed by the enzyme epidermal growth factor receptor kinase. The peptide substrate (RR-SRC) has the sequence arg-arg-leu-ile-glu-asg-ala-glu-tyr-ala-ala-arg-gly. The enzyme used

35 in this assay is the His-tagged cytoplasmic domain of EGFR. A recombinant baculovirus (vHcEGFR52) was constructed containing the EGFR cDNA encoding amino acids 645 - 1186 preceded by Met-Ala-(His)₆. Sf9 cells in 100 mm plates were infected at an moi of 10 pfu/cell and cells were harvested 48 h post infection. A

40 cytoplasmic extract was prepared using 1% Triton X-100 and applied to Ni-NTA column. After washing the column with 20 mM imidazole, HcEGFR was eluted with 250 mM imidazole (in 50 mM Na₂HPO₄, pH 8.0, 300 mM NaCl). Fractions

45 collected were dialyzed against 10 mM HEPES, pH 7.0, 50 mM NaCl, 10% glycerol, 1 µg/mL antipain and leupeptin and 0.1 mM Pefabloc SC. The protein was frozen in dry ice/methanol and stored -70°C.

50

30

55

- 56 -

Test compounds were made into 10 mg/mL stock solutions in 100% dimethylsulfoxide (DMSO). Prior to experiment, stock solutions were diluted to 500 uM with 100% DMSO and then serially diluted to the desired concentration with HEPES buffer (30 mM HEPES pH 7.4).

For the enzyme reaction, 10 μ L of each inhibitor (at various concentrations) were added to each well of a 96-well plate. To this was added 3 μ L of enzyme (1:10 dilution in 10mM HEPES, pH 7.4 for final conc. of 1:120). This was allowed to sit for 10 min on ice and was followed by the addition of 5 μ L peptide (80 μ M final conc.), 10 μ L of 4X Buffer (Table A), 0.25 μ L 33 P-ATP and 12 μ L H₂O. The reaction was allowed to run for 90 min at room temperature and was followed by spotting the entire volume on to precut P81 filter papers. The filter discs were washed 2X with 0.5% phosphoric acid and radioactivity was measured using a liquid scintillation counter.

Table A

Reagent	Final	100 Rxns
1 M HEPES (pH 7.4)	12.5 mM	50 μ L
10mM Na ₃ VO ₄	50 uM	20 μ L
1M MnCl ₂	10 mM	40 μ L
1mM ATP	20 uM	80 μ L
33 P-ATP	2.5uCi	25 μ L

5

- 57 -

10

15

The inhibition data for representative compounds of the invention are shown below in TABLE 1. The IC₅₀ is the concentration of test compound needed to reduce the total amount of phosphorylated substrate by 50%. The % inhibition of the test compound was determined for at least three different concentrations and the IC₅₀ value was evaluated from the dose response curve. The % inhibition was evaluated with the following formula:

20

$$\% \text{ inhibition} = 100 - [\text{CPM}(\text{drug})/\text{CPM}(\text{control})] \times 100$$

25

30

35

where CPM(drug) is in units of counts per minute and is a number expressing the amount of radiolabeled ATP (g-³³P) incorporated onto the RR-SRC peptide substrate by the enzyme after 90 minutes at room temperature in the presence of test compound as measured by liquid scintillation counting. CPM(control) is in units of counts per minute and was a number expressing the amount of radiolabeled ATP (g-³³P) incorporated into the RR-SRC peptide substrate by the enzyme after 90 minutes at room temperature in the absence of test compound as measured by liquid scintillation counting. The CPM values were corrected for the background counts produced by ATP in the absence of the enzymatic reaction. The IC₅₀ values in TABLE 1 are averages of the individual determinations.

20

40

45

50

55

- 58 -

TABLE 1 (recombinant enzyme)
Inhibition of Epidermal Growth Factor Receptor Kinase

<u>Compound</u>	<u>IC₅₀</u> <u>(μM)</u>	<u>Number of</u> <u>Tests</u>
72	0.006	1
74	0.01	1
75	0.0004	2
76	0.01	2
77	0.006	1
79	0.00036	3
82	0.05	1
93	1.0	1
95	0.005	1
96	0.1	1
108	0.026	2
106	0.013	2
107	0.5	1
109	0.007	2
89	0.01	1
115	0.005	1
91	0.015	1
119	0.00005	1
103	0.008	2

Inhibition of Epithelial Cell Kinase (ECK)

In this standard pharmacological test procedure, a biotinylated peptide substrate is first immobilized on neutravidin-coated microtiter plates. The test drug, the Epithelial Cell Kinase (ECK), Mg^{++} , sodium vanadate (a protein tyrosine phosphatase inhibitor), and an appropriate buffer to maintain pH (7.2) are then added

- 59 -

to the immobilized substrate-containing microtiter wells. ATP is then added to initiate phosphorylation. After incubation, the assay plates are washed with a suitable buffer leaving behind phosphorylated peptide which is exposed to horseradish peroxidase (HRP)-conjugated anti-phosphotyrosine monoclonal antibody. The antibody-treated plates are washed again and the HRP activity in individual wells is quantified as a reflection of degree of substrate phosphorylation. This nonradioactive format was used to identify inhibitors of ECK tyrosine kinase activity where the IC_{50} is the concentration of drug that inhibits substrate phosphorylation by 50%. The results obtained for representative compounds of this invention are listed in TABLE 2. Multiple entries for a given compound indicate it was tested multiple times.

Inhibition of Kinase insert Domain containing Receptor (KDR: the catalytic domain of the VEGF receptor)

In this standard pharmacological test procedure, KDR protein is mixed, in the presence or absence of an inhibitor compound, with a substrate peptide to be phosphorylated (a copolymer of glutamic acid and tyrosine, E:Y :: 4:1) and other cofactors such as Mg^{++} and sodium vanadate (a protein tyrosine phosphatase inhibitor) in an appropriate buffer to maintain pH (7.2). ATP and a radioactive tracer (either P^{32} - or P^{33} - labeled ATP) is then added to initiate phosphorylation. After incubation, the radioactive phosphate associated with the acid-insoluble fraction of the assay mixture is then quantified as reflection of substrate phosphorylation. This radioactive format was used to identify inhibitors of KDR tyrosine kinase activity where the IC_{50} is the concentration of drug that inhibits substrate phosphorylation by 50%. The results obtained for representative compounds of this invention are listed in TABLE 2. Multiple entries for a given compound indicate it was tested multiple times.

Mitogen Activated Protein Kinase (MAPK) Assay

To evaluate inhibitors of the MAP (mitogen activated protein) kinase a two component coupled standard pharmacological test procedure, which measures phosphorylation of a serine/threonine residue in an appropriate sequence in the

5

- 60 -

10

15

substrate in the presence and absence of a putative inhibitor, was used. Recombinant human MEK 1 (MAPKK) was first used to activate recombinant human ERK2 (MAPK) and the activated MAPK (ERK) was incubated with substrate (MBP peptide or MYC peptide) in the presence of ATP, Mg^{+2} and radiolabeled ^{33}P ATP. The phosphorylated peptide was captured on a P 81 phosphocellulose filter (paper filter or embedded in microtiter plate) washed and counted by scintillation methods.

20

25

The peptide substrates used in the assay are MBP, peptide substrate (APRTPGGRR), or synthetic Myc substrate, (KKFELLPTPLSPSR•5 TFA. The recombinant enzymes used were prepared as GST fusion proteins of human ERK 2 and human MEK 1. Inhibitor samples were prepared as 10X stocks in 10% DMSO and an appropriate aliquot was used to deliver either 10 ug/ml for a single point screening dose or 100, 10, 1, and 0.1 uM final concentration for a dose response curve. Final DMSO concentrations were less than or equal to 1%.

30

35

40

45

50

The reaction was run as follows in 50 mM Tris kinase buffer, pH 7.4 in a reaction volume of 50 ul. The appropriate volume of kinase buffer and inhibitor sample was added to the tube. Appropriate dilution of enzyme was delivered to give 2-5 ug recombinant MAPK (Erk) per tube. The inhibitor was incubated with MAPK (Erk) for 30 min at 0 deg. C. Recombinant Mek (MAPKK) (0.5-2.5 ug) or fully activated Mek (0.05-0.1 units) was added to activate the Erk and incubated for 30 min at 30° C. Then substrate and gamma ^{33}P ATP was were added to give a final concentration of 0.5- 1 mM MBPP or 250-500 uM Myc; 0.2-0.5 uCi gamma P 33 ATP/tube; 50 μ M ATP final concentration. Samples were incubated at 30°C for 30 minutes and the reaction was stopped by adding 25 μ l of ice cold 10 %TCA .After samples were chilled on ice for 30 min, 20 μ l of sample was transferred onto P 81 phosphocellulose filter paper or appropriate MTP with embedded P 81 filter. Filter papers or MTP were washed 2 times with a large volume of 1% acetic acid, then 2 times with water. The filters or MTP were briefly air dried before addition of scintillant and samples were counted in the appropriate scintillation counter set up for reading ^{33}P isotope. Samples included a positive control (activated enzyme plus substrate); a no enzyme control; a no substrate control; samples with different

55

- 61 -

concentrations of putative inhibitor; and samples with reference inhibitors (other active compounds or non-specific inhibitors such as staurosporine or K252 B).

The raw data was captured as cpm. Sample replicates were averaged and corrected for background count. Mean cpm data was tabulated by group and % inhibition by a test compound was calculated as (corrected cpm control- corrected cpm sample/control) X 100 = % inhibition. If several concentrations of inhibitor were tested, IC_{50} values (the concentration which gives 50% inhibition) were determined graphically from the dose response curve for % inhibition or by an appropriate computer program. The results obtained for representative compounds of this invention are listed in TABLE 2 where there may be separate entries for the same compound; this is an indication that the compound was evaluated more than one time.

TABLE 2

Inhibition of Kinase insert Domain containing Receptor (KDR), Epithelial Cell Kinase (Eck), and Mitogen Activated Protein Kinase (Mek-Erk)

Example	VEGF μ M	Eck μ M	Mek and Erk μ M	erbB2 % Inh 2 μ g/mL
72		> 41.724	> 100	96
74		> 42.982	> 100	95
75		> 37.284	100	
76		> 40.617	> 100	96
77		42.162	> 100	105
78		> 21.269	40	53
79		> 38.610	80	
82	> 1.8315	> 1.832	30	
85		> 21.584	10	87
			4	
			1.8	
			1.8	

Inhibition of Cancer Cell Growth as Measured by Cell Number

Human tumor cell lines were plated in 96-well plates (250 μ l/well, $1-6 \times 10^4$ cells/ml) in RPMI 1640 medium, containing 5% FBS (Fetal Bovine Serum). Twenty four hours after plating, test compounds were added at five log concentrations (0.01-100 mg/ml) or at lower concentrations for the more potent compounds. After 48 hours exposure to test compounds, cells were fixed with trichloroacetic acid, and

- 62 -

stained with Sulforhodamine B. After washing with trichloroacetic acid, bound dye was solubilized in 10 mM Tris base and optical density was determined using plate reader. Under conditions of the assay the optical density is proportional to the number of cells in the well. IC_{50} s (concentrations causing 50% inhibition of cell growth) were determined from the growth inhibition plots. The test procedure is described in details by Philip Skehan et. al, *J.Natl. Canc. Inst.*, 82, 1107-1112 (1990). These data are shown below in TABLE 3 . Information about some of the cell lines used in these test procedures is available from the American Type Tissue Collection: Cell Lines and Hybridomas, 1994 Reference Guide, 8th Edition.

Table 3

Inhibition of Cancer Cell Growth as Measured by Cell Number (IC_{50} μ g/mL)

Example	MDAMB435	SW620	A431	SKBR3	3T3	Her2/3T3
85	27.6	17.98	4.91	1.74		
75	0.2	0.5	0.1	0.04		
74	3.9	0.6	0.9	1.0		
72	3.5	0.4	0.9	0.8		
77	0.1	0.1	0.03	0.02		
79	0.3	0.1	0.04	0.01		
76	0.284	0.239	0.050	0.031		
78	3.194	>5	0.369	1.495		
91	1.83	1.73	0.232	0.181	2.87	0.375
89	2.07	1.53	0.245	0.107	2.04	0.192
90	1.74	1.24	0.234	0.148	2.1	0.329
92	3.32	2.51	0.283	0.188	2.79	0.35

In Vivo Inhibition of the Growth of Human Epidermoid Tumors (A431)

BALB/c nu/nu female mice (Charles River, Wilmington, MA) were used in the in vivo standard pharmacological test procedures. Human epidermoid carcinoma cells A-431 (American Type Culture Collection, Rockville, Maryland # CRL-155) were grown in vitro as described above. A unit of 5×10^6 cells were injected SC

- 63 -

into mice. When tumors attained a mass of between 100 and 150 mg, the mice were randomized into treatment groups (day zero). Mice were treated IP or PO once a day either on days 1, 5, and 9 or on days 1 through 10 post staging with doses of either 80, 40 or 20, or 10 mg/kg/dose of the compound to be evaluated in 0.2% Klucel. Control animals received no drug. Tumor mass was determined every 7 days [(length X width²)/2] for 28 days post staging. Relative tumor growth (Mean tumor mass on day 7, 14, 21, and 28 divided by the mean tumor mass on day zero) is determined for each treatment group. The %T/C (Tumor/Control) is determined by dividing the relative tumor growth of the treated group by the relative tumor growth of the placebo group and multiplying by 100. A compound is considered to be active if the %T/C is found to be significantly less than 100%.

The ability of the compound of Example 92 to inhibit the growth of human epidermoid tumors (A431) in vivo demonstrated below in TABLE 4 below.

TABLE 4
In Vivo Inhibition of the Growth of Human Epidermoid Tumors (A431) in Mice
by the Compound of Example 92

a	b	c,d	b	c,d	b	c,d	b	c,d	e
Drug Treatment mg/kg/dose	Day 7	% T/C	Day 14	% T/C	Day 20	% T/C	Day 28	% T/C	S/T
0.5% Methocel 0.4% Tween 80	5.51		10.43		12.36		14.18		10/10
Example 92 (40 PO)	1.49	27*	1.58	15*	2.60	21*	6.22	44	5/5
Example 92 (10 PO)	3.94	72	10.41	100	14.76	119	22.51	159	5/5

a) compound administered on days 1 through 10 PO.

b) Relative Tumor Growth = $\frac{\text{Mean Tumor Mass on Day 7, 14, 21, 28}}{\text{Mean Tumor Mass on Day 0}}$

c) % T/C = $\frac{\text{Relative Tumor Growth of Treated Group}}{\text{Relative Tumor Growth of Placebo Group}} \times 100$

d) Statistical Analysis (Student's T-test) of Log Relative Tumor Growth. * Indicates statistically ($p < 0.01$) significant reduction in Relative Tumor Growth of Treated Group compared to the Placebo Control.

e) S/T = # of Survivors/# of Treated on Day +28 post tumor staging.

- 64 -

As indicated by the results presented in TABLE 4, the compound of Example 92 is an effective inhibitor of tumor growth in vivo when given orally at 40 mg/Kg.

The ability of the compound of Example 89 to inhibit the growth of human epidermoid tumors (A431) in vivo demonstrated below in TABLE 5 below.

TABLE 5
In Vivo Inhibition of the Growth of Human Epidermoid Tumors (A431) in Mice
by the Compound of Example 89

a	b	c,d	b	c,d	b	c,d	b	c,d	e
Drug Treatment mg/kg/dose	Day 7	% T/C	Day 14	% T/C	Day 21	% T/C	Day 28	% T/C	S/T
0.5% Methocel 0.4% Tween 80	4.18		10.44		15.08		28.23		9/10
Example 89 (40 PO)	0.49	11*	0.58	6*	3.11	21*	7.20	26*	5/5
Example 89 (10 PO)	2.09	50*	3.37	32*	5.76	38*	7.24	26*	4/5

a) compound administered on days 1 through 10 PO.

b) Relative Tumor Growth = $\frac{\text{Mean Tumor Mass on Day 7, 14, 21, 28}}{\text{Mean Tumor Mass on Day 0}}$

c) % T/C = $\frac{\text{Relative Tumor Growth of Treated Group}}{\text{Relative Tumor Growth of Placebo Group}} \times 100$

d) Statistical Analysis (Student's T-test) of Log Relative Tumor Growth. * Indicates statistically ($p < 0.01$) significant reduction in Relative Tumor Growth of Treated Group compared to the Placebo Control.

e) S/T = # of Survivors/# of Treated on Day +28 post tumor staging.

As indicated by the results presented in TABLE 5, the compound of Example 89 is an effective inhibitor of tumor growth in vivo when given orally at 40 mg/Kg and 10 mg/Kg.

Based on the results obtained for representative compounds of this invention, the compounds of this invention are antineoplastic agents which are useful in treating, inhibiting the growth of, or eradicating neoplasms. In particular, the compounds of this invention are useful in treating, inhibiting the growth of, or eradicating neoplasms that express EGFR such as those of the breast, kidney, bladder, mouth, larynx, esophagus, stomach, colon, ovary, or lung. In addition, the compounds of this invention are useful in treating, inhibiting the growth of, or eradicating neoplasms of

5

- 65 -

10

the breast that express the receptor protein produced by the erbB2 (Her2) oncogene. Based on the results obtained, the compounds of this invention are also useful in the treatment of polycystic kidney disease.

15

5 The compounds of this invention may formulated neat or may be combined with one or more pharmaceutically acceptable carriers for administration. For example, solvents, diluents and the like, and may be administered orally in such forms as tablets, capsules, dispersible powders, granules, or suspensions containing, for example, from about 0.05 to 5% of suspending agent, syrups containing, for example, from about 10 to 50% of sugar, and elixirs containing, for example, from about 20 to 50% ethanol, and the like, or parenterally in the form of sterile injectable solution or suspension containing from about 0.05 to 5% suspending agent in an isotonic medium. Such pharmaceutical preparations may contain, for example, from about 0.05 up to about 90% of the active ingredient in combination with the carrier, 15 more usually between about 5% and 60% by weight.

30

The effective dosage of active ingredient employed may vary depending on the particular compound employed, the mode of administration and the severity of the condition being treated. However, in general, satisfactory results are obtained when the compounds of the invention are administered at a daily dosage of from about 0.5 35 to about 1000 mg/kg of body weight, optionally given in divided doses two to four times a day, or in sustained release form. The total daily dosage is projected to be from about 1 to 1000 mg, preferably from about 2 to 500 mg. Dosage forms suitable for internal use comprise from about 0.5 to 1000 mg of the active compound in intimate admixture with a solid or liquid pharmaceutically acceptable carrier. This 40 dosage regimen may be adjusted to provide the optimal therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

45

50

The compounds of this invention may be administered orally as well as by intravenous, intramuscular, or subcutaneous routes. Solid carriers include starch, lactose, dicalcium phosphate, microcrystalline cellulose, sucrose and kaolin, while 30 liquid carriers include sterile water, polyethylene glycols, non-ionic surfactants and

55

5

- 66 -

10

edible oils such as corn, peanut and sesame oils, as are appropriate to the nature of the active ingredient and the particular form of administration desired. Adjuvants customarily employed in the preparation of pharmaceutical compositions may be advantageously included, such as flavoring agents, coloring agents, preserving agents, and antioxidants, for example, vitamin E, ascorbic acid, BHT and BHA.

15

The preferred pharmaceutical compositions from the standpoint of ease of preparation and administration are solid compositions, particularly tablets and hard-filled or liquid-filled capsules. Oral administration of the compounds is preferred.

20

In some cases it may be desirable to administer the compounds directly to the airways in the form of an aerosol.

25

The compounds of this invention may also be administered parenterally or intraperitoneally. Solutions or suspensions of these active compounds as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxy-propylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

30

35

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

40

45

For the treatment of cancer, the compounds of this invention can be administered in combination with other antitumor substances or with radiation therapy. These other substances or radiation treatments can be given at the same or at different times as the compounds of this invention. These combined therapies may effect synergy and result in improved efficacy. For example, the compounds of this

50

55

- 67 -

invention can be used in combination with mitotic inhibitors such as taxol or vinblastine, alkylating agents such as cisplatin or cyclophosphamide, antimetabolites such as 5-fluorouracil or hydroxyurea, DNA intercalators such as adriamycin or bleomycin, topoisomerase inhibitors such as etoposide or camptothecin, antiangiogenic agents such as angiostatin, and antiestrogens such as tamoxifen.

The preparation of representative examples of the compounds of this invention is described below.

Example 1

1,4-Dihydro-7-methoxy-4-oxo-quinoline-3-carbonitrile

A mixture of 30.2 g (245.2 mmol) of 3-methoxy aniline and 41.5 g (245.2 mmol) of ethyl(ethoxymethylene) cyanoacetate was heated in the absence of solvent to 140°C for 30 minutes. To the resulting oil was added 1200 ml of Dowtherm. The solution was refluxed with stirring under nitrogen for 22 hours. The mixture was cooled to room temperature and solid was collected and washed with hexanes. The solid was recrystallized from acetic acid to give 17 g of 1,4-dihydro-7-methoxy-4-oxo-quinoline-3-carbonitrile: mass spectrum (electrospray, m/e): M+H 200.9.

Example 2

1,4-Dihydro-7-methoxy-6-nitro-4-oxo-quinoline-3-carbonitrile

To a suspension of 10 g (49.6 mmol) of 1,4-dihydro-7-methoxy-4-oxo-quinoline-3-carbonitrile in 160 ml of trifluoroacetic anhydride was added 6 g (74.9 mmol) of ammonium nitrate over a period of 3 hours. The mixture was stirred an additional two hours. Excess anhydride was removed at reduced pressure at 45°C. The residue was stirred with 500 ml of water. The solid was collected and washed with water. The solid was dissolved in 1000 ml of boiling acetic acid and the solution was treated with decolorizing charcoal. The mixture was filtered and concentrated to a volume of 300 ml. Cooling gave a solid which was collected giving 5.4 g of 1,4-dihydro-7-methoxy-6-nitro-4-oxo-quinoline-3-carbonitrile as a brown solid: mass spectrum (electrospray, m/e): M+H 246.

5

- 68 -

Example 3

10

4-Chloro-7-methoxy-6-nitro-quinoline-3-carbonitrile

15

A mixture of 5.3 g (21.6 mmol) of 1,4-dihydro-7-methoxy-6-nitro-4-oxo-quinoline-3-carbonitrile and 9 g (43.2 mmol) of phosphorous pentachloride was heated at 165°C for 2 hours. The mixture was diluted with hexanes and the solid was collected. The solid was dissolved in 700 ml ethyl acetate and washed with cold dilute sodium hydroxide solution. The solution was dried over magnesium sulfate and filtered through a pad of silica gel giving 5.2 g of 4-chloro-7-methoxy-6-nitro-quinoline-3-carbonitrile as a tan solid.

20

10

Example 4

25

4-[(3-Bromophenyl)amino]-7-methoxy-6-nitro-quinoline-3-carbonitrile

30

A solution of 5.2 g (19.7 mmol) of 4-chloro-7-methoxy-6-nitro-quinoline-3-carbonitrile and 3.7 g (21.7 mmol) of 3-bromo aniline in 130 ml of methoxyethanol was refluxed under nitrogen for 4 hours. The reaction mixture was poured into dilute sodium bicarbonate solution. Solid was collected and washed with water and dried in air. The solid was chromatographed on silica gel eluting with chloroform-ethyl acetate 9:1. Solvent was removed from product fractions giving 1.2 g of 4-[(3-bromophenyl)amino]-7-methoxy-6-nitro-quinoline-3-carbonitrile as a yellow solid:

35

mass spectrum (electrospray, m/e): M+H 399.0, 402.0.

20

Example 5

40

6-Amino-4-[(3-bromophenyl)amino]-7-methoxy-quinoline-3-carbonitrile

45

A mixture of 2.05 g (5.1 mmol) of 4-[(3-bromophenyl)amino]-7-methoxy-6-nitro-quinoline-3-carbonitrile, 1.37 g (25.7 mmol) of ammonium chloride, and 0.86 g (15.4 mmol) of powdered iron was stirred at reflux in 26 ml water and 26 ml methanol for 2 hours. The mixture was diluted with ethyl acetate and the hot mixture was filtered. The organic layer was separated from the filtrate and dried over magnesium sulfate. The solvent was removed and the residue was chromatographed on silica gel eluting with mixtures of chloroform and ethyl acetate. Product fractions were combined to

50

55

- 69 -

give 1.3 g of 6-amino-4-[(3-bromophenyl)amino]-7-methoxy-quinoline-3-carbonitrile as a yellow solid: mass spectrum (electrospray, m/e): M+H 369.1, 371.1.

Example 6

2-Cyano-3-(4-nitrophenylamino)acrylic Acid Ethyl Ester

4-Nitroaniline (60.0g, 0.435mol) and ethyl(ethoxymethylene) cyanoacetate (73.5g, 0.435mol) were mixed mechanically in a flask. The mixture was heated at 100°C for 0.5h after it had melted and resolidified. A 114 g portion of the crude product was recrystallized from dimethylformamide to give 44.2g of yellow crystals; mp 227-228.5°C.

Example 7

1,4-Dihydroquinoline-6-Nitro-4-oxo-3-carbonitrile

A slurry of 25.0g (95.8mmol) of 2-cyano-3-(4-nitrophenylamino)acrylic acid ethyl ester in 1.0L of Dowtherm A was heated at 260°C under N₂ for 12.5h. The cooled reaction was poured into 1.5L of hexane. The product was collected, washed with hexane and hot ethanol and dried in vacuo. There was obtained 18.7g of brown solid. An analytical sample was obtained by recrystallization from dimethylformamide/ethanol: mass spectrum (electrospray, m/e): M+H 216.

Example 8

4-Chloro-6-nitro-quinoline-3-carbonitrile

A mixture of 31.3g (0.147mol) of 6-nitro-4-oxo-1,4-dihydro-quinoline-3-carbonitrile and 160mL of phosphorous oxychloride was refluxed for 5.5h. The phosphorous oxychloride was removed in vacuo and the residue was poured over ice and neutralized with sodium bicarbonate. The product was collected, washed with water and dried in vacuo (50°C). There was obtained 33.5g of tan solid; solid: mass spectrum (electrospray, m/e): M+H 234.

- 70 -

Example 9**4-[3-(3-Bromophenyl)amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 17.0g (73.1mmol) of 4-chloro-6-nitro-quinoline-3-carbonitrile and 15.1g (87.7mmol) of 3-bromoaniline in 425mL of ethanol was refluxed for 5h.

Saturated sodium bicarbonate was added and then all volatile material was removed in vacuo. The residue was slurried with hexane and the product was collected and washed with hexane. The crude product was washed with water and dried in vacuo(60°C). There was obtained 22.5g of yellow solid. An analytical sample was obtained by recrystallization from ethyl acetate; mp 258-259°C.

Example 10**6-Amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 4.00g (10.8mmol) of 4-[(3-bromophenyl)amino]-6-nitro-quinoline-3-carbonitrile and 12.2g (54.2mmol) of SnCl₂ dihydrate in 160mL of ethanol was refluxed under N₂ for 1.3h. After cooling to 25°C, ice water and sodium bicarbonate were added and the mixture was stirred for 2h. Extraction with chloroform, treatment with Darco, drying (magnesium sulfate) and solvent removal gave 3.9g of brown crystals: mass spectrum (electrospray, m/e): M+H 339.

Example 11**4-[3,4-Dibromophenyl]amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 6.20g (26.6mmol) of 4-chloro-6-nitro-quinoline-3-carbonitrile and 8.00g (31.9mmol) of 3,4-dibromoaniline in 160mL of ethanol was refluxed under N₂ for 5h. Saturated sodium bicarbonate was added and volatile material was removed. The residue was slurried with hexane, collected, washed with hexane and water and dried. The insoluble material was repeatedly extracted with boiling ethyl acetate and the solution was then filtered through silica gel. The solvent was removed to give 3.80g of green solid: mass spectrum (electrospray, m/e): M+H 449.

- 71 -

Example 12**6-Amino-4-[(3,4-dibromophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 4.90g (10.9mmol) of 4-[(3,4-dibromophenyl)amino]-6-nitro-quinoline-3-carbonitrile and 12.4g (54.7mmol) of SnCl₂ dihydrate in 200mL of ethanol was refluxed under N₂ for 1.5h. After cooling to 25°C, the reaction was diluted with ice water, neutralized with sodium bicarbonate and stirred for 2h. This solution was then extracted with chloroform, treated with Darco, dried(magnesium sulfate) and evaporated. After drying in vacuo(40°C), there was obtained 1.25g of brown solid: mass spectrum (electrospray, m/e): M+H 417, 419, 421.

Example 13**6-Nitro-4-[(3-trifluoromethylphenyl)amino]-quinoline-3-carbonitrile**

A mixture of 10.6g (45.7mmol) of 4-chloro-6-nitro-quinoline-3-carbonitrile and 8.82g (54.8mmol) of 3-(trifluoromethyl)aniline in 270 mL of ethanol was refluxed under N₂ for 5h. The reaction was diluted with ethanol, neutralized with satd sodium bicarbonate and evaporated. The residue was slurried with hexane, collected, washed with hexane and water and dried in vacuo(60°C) to give 10.9g of yellow solid. A 2.00g sample was recrystallized from ethanol to give 1.20g of bright yellow solid; mp 260-261°C.

Example 14**6-Amino-4-[(3-trifluoromethylphenyl)amino]-quinoline-3-carbonitrile**

A slurry of 6.00g(16.8mmol) of 6-nitro-4-[(3-trifluoromethylphenyl)amino]-quinoline-3-carbonitrile and 18.9g (83.3mmol) of SnCl₂ dihydrate in 240 mL of ethanol was refluxed under N₂ for 1h. After cooling to 25°C, the reaction was diluted with ice water, neutralized with sodium bicarbonate and stirred for 2h. The product was extracted with chloroform, treated with Darco, dried(magnesium sulfate) and evaporated. The residue was filtered through silica gel(10% methanol in chloroform),

5

- 72 -

10

evaporated and dried in vacuo(40°C) to give 4.87g of brown solid: mass spectrum (electrospray, m/e): M+H 329.

Example 15

15

5 4-Bromo-but-2-enoic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide.

20

A solution of 1.65 grams (0.01 mole) of 4-bromo crotonic acid (Giza Braun, J. Am. Chem. Soc. 52, 3167 1930) in 15ml of dichloromethane was treated with 1.74 ml (0.02 moles) of oxalyl chloride and 1 drop of N, N- dimethylformamide. After an hour the solvents were removed on the rotary evaporator. The 4-bromo crotonyl chloride was taken up in 25 ml of tetrahydrofuran, and 3.39 grams of 6-Amino-4-(3-bromo-phenylamino)-quinoline-3-carbonitrile in 25 ml of tetrahydrofuran was added dropwise. This was followed by the dropwise addition of 1.92 ml (0.011 moles) of diisopropylethylamine. After the addition of 25 ml of water and 50 ml of ethyl acetate, the layers were separated. The organic layer was dried over anhydrous sodium sulfate, and taken to a solid *in vacuo*. This solid was digested for an hour with refluxing ethyl acetate then filtered from the ethyl acetate while still hot. Thus was obtained 3.31 grams (68%) of 4-bromo-but-2-enoic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide.

25

30

35

20

Example 16

2-Cyano-3-(2-methyl-4-nitrophenyl)acrylic Acid Ethyl Ester

40

A mixture of 2-methyl-4-nitroaniline (38.0 g, 250 mmol), ethyl (ethoxymethylene)-cyanoacetate (50.8 g, 300 mmol), and 200 ml of toluene was refluxed for 24 h, cooled, diluted with 1:1 ether-hexane, and filtered. The resulting white solid was washed with hexane-ether and dried to give 63.9 g, mp 180-210°C.

45

Example 17

1,4-Dihydroquinoline-8-methyl-6-nitro-3-carbonitrile

50

30 A stirred mixture of 64 g (230 mmol) of 2-cyano-3-(2-methyl-4-nitrophenyl)acrylic acid ethyl ester and 1.5 L of Dowtherm A was heated at 260°C for 12 h, cooled,

55

5

- 73 -

10

diluted with hexane, and filtered. The grey solid thus obtained was washed with hexane and dried to give 51.5 g, mp 295-305°C.

Example 18

15

5 4-Chloro-8-methyl-6-nitro-quinoline-3-carbonitrile

20

A stirred mixture of 1,4-dihydroquinoline-8-methyl-6-nitro-3-carbonitrile (47 g, 200 mmol) and 200 ml of phosphorous oxychloride was refluxed for 4 h. The phosphorous oxychloride was removed *in vacuo*, and the residue was stirred with methylene chloride at 0°C and treated with a slurry of ice and sodium carbonate. The organic layer was separated and washed with water. The solution was dried and concentrated to a volume of 700 ml. The product was precipitated by the addition of hexane and cooling to 0°C. The white solid was filtered off and dried to give 41.6 g, mp 210-212°C.

25

Example 19

30

15 4-[(3-Bromophenyl)amino]-8-methyl-6-nitro-quinoline-3-carbonitrile

35

A stirred mixture of 4-chloro-8-methyl-6-nitro-quinoline-3-carbonitrile (14.8 g, 60 mmol), 3-bromoaniline (12.4 g, 72 mmol), pyridine hydrochloride (6.93 g, 60 mmol), and 180 ml of ethoxyethanol was refluxed for 1.5 h, cooled, poured into a stirred mixture of water and an amount of sodium carbonate to give a pH of 8-9. The resulting yellow solid was filtered, washed with water, dried, digested in boiling ether, filtered, and dried to give 22.6 g, mp 263-267°C.

20

40

Example 20

45

4-[(3-Bromophenyl)-N-acetylamino]-8-methyl-6-nitro-quinoline-3-carbonitrile

50

25 A stirred mixture of 4-[(3-bromophenyl)amino]-8-methyl-6-nitro-quinoline-3-carbonitrile (15.3 g, 40 mmol), 0.37 g (3 mmol) of dimethylaminopyridine, 40 ml of acetic anhydride, and 80 ml of pyridine was refluxed for 3 h and concentrated at 50°C under vacuum. The residue was stirred with methylene chloride and 0.1 N HCl. After filtration through Celite, the organic layer was washed with water, dried and concentrated. The residue was subjected to chromatography on silica gel with 1%

55

5

- 74 -

10

acetic acid in methylene chloride to give 11.2 g of an amber glass, NMR (CDCl₃) d 2.29 (N-acetyl group).

Example 21

15

5 8-Bromomethyl-4-[(3-bromophenyl)-N-acetylamino]-6-nitro-quinoline-3-carbonitrile

20

A stirred mixture of 4-[(3-bromophenyl)-N-acetylamino]-8-methyl-6-nitro-quinoline-3-carbonitrile (10.6 g, 25 mmol), N-bromosuccinimide (6.68 g, 37.5 mmol), 0.30 g of dibenzoyl peroxide, and 200 ml of carbon tetrachloride was refluxed for 2h, treated with an additional 0.30 g of dibenzoyl peroxide, and refluxed an additional 2.5 h, cooled, diluted with methylene chloride, and stirred with aqueous sodium bisulfite. The organic layer was separated and washed successively with water, sodium bicarbonate solution, and water. The solution was dried and

25 evaporated to give 15 g of a white foam, NMR (CDCl₃) d 5.19 (dd, CH₂Br).

15

Example 22

30

4-[(3-Bromophenyl)amino]-8-dimethylaminomethyl-6-nitro-quinoline-3-carbonitrile

35

To a stirred solution of dimethylamine in THF (2.0 M; 115 ml; 230 mmol) at 0°C was added a solution of 8-bromomethyl-4-[(3-bromophenyl)-N-acetylamino]-6-nitro-quinoline-3-carbonitrile (11.6 g, 23 mmol) in 115 ml of THF during 15 m. After

20 warming to 25°C the mixture was stirred for 2h. The THF was evaporated off, and the residue was refluxed in 230ml of methanol with 12.7 g (92 mmol) of potassium carbonate for 1 h. The mixture was cooled, saturated with CO₂, and concentrated. The residue was partitioned with methylene chloride and water. The organic layer was washed with water, dried, and concentrated. The residue was subjected to

25 chromatography on silica gel with methylene chloride-ethyl acetate-methanol-triethylamine to give 6.0 g yellow solid, mp 223-226°C.

45

50

55

5

- 75 -

Example 23

10

6-Amino-4-[(3-bromophenyl)amino]-8-dimethylaminomethyl-quinoline-3-carbonitrile

15

20

25

A stirred mixture of 4-[(3-bromophenyl)amino]-8-dimethylaminomethyl-6-nitro-quinoline-3-carbonitrile (5.98 g, 14.1 mmol), iron powder (2.76 g, 49 mg-atoms), acetic acid (5.67 ml, 99 mmol), and 70 ml of methanol was refluxed for 2 h and then evaporated to remove methanol. The residue was stirred with water for 10 m, and the orange solid was filtered off and washed with 2% acetic acid. The total filtrate was basified to pH 10 with 5 N sodium hydroxide. The resulting precipitate was extracted with methylene chloride. The extract was washed with water, dried, and concentrated. The residue was subjected to chromatography on silica gel with ethyl acetate-methanol-triethylamine to give 3.34 g of amber solid; mass spectrum (electrospray, m/e) M+H 396.2, 398.1.

15

Example 24

30

6-Amino-4-[(3-iodophenyl)amino]-quinoline-3-carbonitrile

35

40

A mixture of 6.70g (16.1 mmol) 4-[(3-iodophenyl)amino]-6-nitro-quinoline-3-carbonitrile, 300 ml ethanol, and 18.2 g (80.5 mmol) SnCl₂ dihydrate was heated to reflux under N₂. Removed heat at 2 hours, added ice water. Added sodium bicarbonate until pH was basic, forming a thick yellow mixture. Stirred for 2 ½ hours. Extracted with chloroform, stirred organic portion with Darco and filtered through magnesium sulfate. Stripped solvent and dried in vacuo, giving 3.48g of yellow-brown solid: mass spectrum (electrospray m/e): M+H = 387.0.

25

Example 25

45

4-[(3-Iodophenyl)amino]-6-nitro-quinoline-3-carbonitrile

50

55

A mixture of 3.10 ml (25.7 mmol) 3-iodoaniline, 200 ml ethanol, and 5.00 g (21.4 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile was heated to reflux under N₂ for 3 ½ hours. Cooled and made basic with a saturated sodium bicarbonate. Stripped solvents and azeotrope with ethanol. Slurried residue with hexane and collected. Air dried, washed solids with water, and dried in vacuo. Dissolved solids in 400 ml

- 76 -

ethyl acetate, stirred with Darco, filtered and removed solvent. Dried solids in vacuo to give 7.38 g of yellow solid: mass spectrum (electrospray m/e): M+H = 417.0.

Example 26**6-Amino-4-[(3-methylphenyl)amino]-quinoline-3-carbonitrile**

Added 253 mg 10% palladium on carbon to a round bottom flask under N₂ and covered catalyst with 140 ml ethanol. To this added 2.49 g (8.18 mmol) 6-nitro-4-[(3-methylphenyl)amino]-quinoline-3-carbonitrile and 640 µl (20.4 mmol) anhydrous hydrazine. The mixture was heated to reflux for 2 hours 15 minutes and filtered hot through celite. Stripped solvent and dried in vacuo, giving 2.455 g of yellow solid: mass spectrum (electrospray m/e): M+H = 275.2.

Example 27**6-Nitro-4-[(3-methylphenyl)amino]-quinoline-3-carbonitrile**

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol, and 2.75 ml (25.7 mmol) 3-toluidine was heated to reflux for 4 ½ hours. Cooled and added a saturated sodium bicarbonate until pH was basic. Stripped solvents and azeotroped with ethanol. Slurried with hexane, collected, and air dried. Washed with water and dried in vacuo. Boiled in ethyl acetate, stirred with Darco and filtered. Stripped solvent and dried in vacuo to give 4.82 g of yellow-orange solid: mass spectrum (electrospray m/e): M+H = 305.2.

Example 28**6-Amino-4-[(3-chlorophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 6.30 g (19.4 mmol) 4-[(3-chlorophenyl)amino]-6-nitro-quinoline-3-carbonitrile, 300 ml ethanol, and 21.9 g (97 mmol) SnCl₂ dihydrate were heated to reflux under N₂. Removed heat at 2 ½ hours, added ice water and made basic with sodium bicarbonate. Stirred for 2 hours and extracted with chloroform. Dried organic layer with sodium sulfate, filtered, stripped solvent and dried residue in vacuo, giving 5.74 g of yellow-brown solid: mass spectrum (electrospray m/e): M+H = 295.1, 297.1.

5

- 77 -

Example 29

10

4-[(3-Chlorophenyl)amino]-6-nitro-quinoline-3-carbonitrile

15

A mixture of 10.0 g (42.9 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 260 ml ethanol, and 5.40 ml 3-chloroaniline was heated to reflux under N₂. Removed heat at 4 hours, cooled to 25°C and added saturated sodium bicarbonate until the pH was basic. Stripped solvents and azeotroped with ethanol. Slurried residue with hexane, collected solid, and air dried. Washed solids with water and dried in vacuo. Dissolved in boiling ethyl acetate, stirred with Darco, and filtered. Stripped solvent and dried residue in vacuo, giving 6.5g of yellow solid: mass spectrum (electrospray m/e): M+H = 325.0, 327.0.

20

10

Example 30

25

6-Amino-4-[(3-methoxyphenyl)amino]-quinoline-3-carbonitrile

30

325 mg of 10% palladium on carbon was added to a round bottom flask under N₂ and covered with 165 ml ethanol. Added 3.29 g (10.3 mmol) 4-[(3-methoxyphenyl)-amino]-6-nitro-quinoline-3-carbonitrile and 800µl anhydrous hydrazine and heated mixture to reflux. At 1 ½ hours, filtered hot through celite, stripped solvent and dried in vacuo, giving 2.876 g of yellow solid: mass spectrum (electrospray m/e): M+H = 291.2.

35

20

Example 31

40

4-[(3-methoxyphenyl)amino]-6-nitro-quinoline-3-carbonitrile

45

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol, and 3.0 ml (26.0 mmol) m-anisidine was heated to reflux under N₂. Removed heat at 4 ½ hours and made basic with saturated sodium bicarbonate. Stripped solvents and azeotroped with ethanol. Slurried with hexane and collected crystals. Washed with water, dried in vacuo. Dissolved 5.94 g of crude product in 320 ml boiling ethyl acetate, stirred with Darco, filtered, stripped solvent, and dried in vacuo, giving about 5 g of yellow-orange solid: mass spectrum (electrospray m/e): M+H = 291.1.

50

30

55

- 78 -

Example 32**6-Amino-4-[(3-chloro-4-fluorophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 5.360 g (15.6 mmol) 4-[(3-chloro-4-fluorophenyl)amino]-6-nitro-quinoline-3-carbonitrile, 250 ml ethanol, and 17.67 g (78.2 mmol) SnCl_2 dihydrate was heated to reflux under N_2 . Removed heat at 1 ½ hours and added ice water. Made basic with sodium bicarbonate. Stirred for 2 hours extracted with chloroform. Added brine to the separatory funnel to help separate layers. Stirred organic layer with Darco and dried with sodium sulfate. Filtered, stripped solvent and dried in vacuo, giving 4.460 g of yellow-brown solid: mass spectrum (electrospray m/e): $\text{M}+\text{H} = 312.9, 315.0$.

Example 33**4-[(3-chloro-4-fluorophenyl)amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol, and 3.75 g (25.8 mmol) 3-chloro-4-fluoroaniline was heated to reflux under N_2 . Removed heat at 3 ½ hours and added a solution of saturated sodium bicarbonate until mixture was basic. Stripped solvents and azeotroped with ethanol. Slurried residue with hexane, collected solids, washed with water and dried in vacuo. Dissolved solids in 250 ml boiling ethyl acetate, stirred with Darco, and filtered. Stripped solvent and dried in vacuo, giving 6.036 g of yellow solid: mass spectrum (electrospray m/e): $\text{M}+\text{H} = 343.1, 345.1$.

Example 34**6-Amino-4-[(4-bromophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 3.10 g (8.40 mmol) 4-[(4-bromophenyl)amino]-6-nitro-quinoline-3-carbonitrile, 155 ml ethanol, and 9.47 g (42.0 mmol) SnCl_2 dihydrate was heated to reflux under N_2 . After 4 hours, removed heat and added ice water. Made basic with sodium bicarbonate and stirred for 2 hours. With mixture still basic, extracted with chloroform, stirred organic layer with Darco and dried with sodium sulfate. Filtered, stripped solvent and dried in vacuo, giving 2.265 g of brown-yellow solid: mass spectrum (electrospray m/e): $\text{M}+\text{H} = 339.0, 341.0$.

- 79 -

Example 35**4-[(4-Bromophenyl)amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol, and 4.42 g (25.8 mmol) p-bromoaniline was heated to reflux under N₂ for 3 hours. Removed heat and added saturated sodium bicarbonate until basic. Stripped solvents and azeotroped with ethanol. Slurried residue with hexane, collected solids, and air dried. Washed with water and dried in vacuo. Boiled in 1.4 liters ethyl acetate, and without completely dissolving all solids, stirred with Darco, and filtered. Stripped solvent and dried in vacuo, giving 3.524 g of yellow solid: mass spectrum (electrospray m/e): M+H = 369, 370.9.

Example 36**6-Amino-4-[(3,4-difluorophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 4.53 g (13.9 mmol) 4-[(3,4-difluorophenyl)amino]-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol and 15.72 g (69.4 mmol) SnCl₂ dihydrate was heated to reflux under N₂. Removed heat at 1 ½ hours, added ice water and made basic with sodium bicarbonate. Stirred for 2 hours and extracted with chloroform. Stirred organic layer with Darco, dried with sodium sulfate and filtered. Stripped solvent and dried in vacuo, giving 3.660 g of yellow-green solid: mass spectrum (electrospray m/e): M+H = 297.1.

Example 37**4-[(3,4-Difluorophenyl)amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 250 ml ethanol and 2.55 ml (25.8 mmol) 3,4-difluoroaniline was heated to reflux under N₂. Removed heat at 3 ½ hours and made basic with saturated sodium bicarbonate. Stripped solvents and azeotroped with ethanol. Slurried residue with hexane, collected solids and air dried. Washed with water and dried in vacuo. Dissolved in ethyl acetate, stirred with Darco, filtered, stripped solvent and dried in vacuo, giving 5.02 g of yellow solid: mass spectrum (electrospray m/e): M+H = 327.1.

- 80 -

Example 38**6-Amino-4-[(3-chloro-4-thiophenoxyphenyl)amino]-quinoline-3-carbonitrile**

A mixture of 6.753 g (15.6 mmol) 4-[(3-chloro-4-thiophenoxyphenyl)amino]-6-nitro-quinoline-3-carbonitrile, 250 ml ethanol, and 17.66 g (78.0 mmol) SnCl_2 dihydrate was heated to reflux under N_2 . Removed heat at 2 hours, added large volume of ice water, and made basic with sodium bicarbonate. Stirred for 2 hours and with mixture still basic, extracted with chloroform. Stirred organic layer with Darco, dried with sodium sulfate, filtered, stripped solvent and dried in vacuo, giving 5.996 g of yellow-brown solid: mass spectrum (electrospray m/e): $\text{M}+\text{H} = 403.1, 405.1$.

Example 39**4-[(3-Chloro-4-thiophenoxyphenyl)amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 250 ml ethanol, and 6.07 g (25.6 mmol) 3-chloro-4-thiophenoxyaniline was heated to reflux under N_2 . Removed heat at about 8 hours, made basic with saturated sodium bicarbonate, stripped solvents and azeotroped with ethanol. Slurried residue with hexane and collected solids. Washed with water and dried in vacuo. Dissolved nearly completely in 400 ml ethyl acetate, stirred with Darco and filtered. Stripped solvent and boiled in hexane to remove last of the excess aniline. Dried in vacuo, giving 6.90 g of red solid: mass spectrum (electrospray m/e): $\text{M}+\text{H} = 433.1, 435.1$.

Example 40**6-Amino-4-[(3-cyanophenyl)amino]-quinoline-3-carbonitrile**

Added 100 mg of 10% palladium on carbon to a round bottom flask under N_2 and covered with 50 ml ethanol. Added 1.00g (3.17 mmol) 4-[(3-cyanophenyl)amino]-6-nitro-quinoline-3-carbonitrile and 250 μl (7.39 mmol) anhydrous hydrazine and heated to reflux. Removed heat at 2 hours and filtered hot through celite. Stripped solvent and dried in vacuo, giving 887 mg of yellow solid: mass spectrum (electrospray m/e): $\text{M}+\text{H} = 286.2$.

5

- 81 -

Example 41

10

4-[(3-Cyanophenyl)amino]-6-nitro-quinoline-3-carbonitrile

15

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol, and 3.04 g (25.8 mmol) 3-aminobenzonitrile was heated to reflux. Removed heat at 3 ½ hours and made basic with saturated sodium bicarbonate. Stripped solvents and air dried. Slurried residue with hexane and collected solids. Washed with water and dried in vacuo. Boiled in large volume ethyl acetate, collected solids and dried in vacuo, giving 5.15 g of yellow-brown solid: mass spectrum (electrospray m/e): 316.0.

20

10

Example 42

25

6-Amino-4-[(3-ethynylphenyl)amino]-quinoline-3-carbonitrile

30

A mixture of 2.00g (6.36 mmol) 4-[(3-ethynylphenyl)amino]-6-nitro-quinoline-3-carbonitrile, 100 ml ethanol, and 7.19 g (31.8 mmol) SnCl₂ dihydrate was heated to reflux under N₂. Removed heat at 3 ½ hours and added ice water. Made basic with sodium bicarbonate and stirred for 2 hours. Extracted with chloroform, stirred organic layer with Darco, dried with sodium sulfate, filtered, stripped solvent, and dried in vacuo, giving 1.737 g of yellow-brown solid: mass spectrum (electrospray m/e): M+H = 285.2.

35

20

Example 43

40

4-[(3-Ethynylphenyl)amino]-6-nitro-quinoline-3-carbonitrile

45

A mixture of 5.00 g (21.5 mmol) 4-chloro-6-nitro-quinoline-3-carbonitrile, 200 ml ethanol, and 3.82 g (32.6 mmol) 3-ethynylaniline was heated to reflux under N₂. Removed heat at 3 ½ hours and added a solution of saturated sodium bicarbonate until basic. Stripped solvents and azeotroped with ethanol. Slurried residue with hexane and collected solids. Washed with water and dried in vacuo. Dissolved in ethyl acetate, stirred with Darco, filtered, stripped solvent and dried in vacuo, giving 4.544 g of yellow solid: mass spectrum (electrospray m/e): M+H = 315.1.

50

30

55

- 82 -

Example 44**4-[(3-Bromo-4-fluorophenyl)amino]-6-nitro-quinoline-3-carbonitrile**

A mixture of 3.8g (16.33mmol) of 4-chloro-6-nitro-quinoline-3-carbonitrile and 3.7g (20mmol) of 3-bromo-4-fluoroaniline in 200mL of ethanol was refluxed for 3hr.

After the solvent was removed, the residue was dissolved in ethyl acetate and washed with sodium bicarbonate. The product was collected as a pale yellow solid, 6.5g (71%); ESMS m/z 387.3, 389.2, mp 269-270°C (dec).

Example 45**6-amino-4-[(3-Bromo-4-fluorophenyl)amino]-quinoline-3-carbonitrile**

A mixture of 8g (20.67mmol) of 4-[(3-chloro-4-fluorophenyl)amino]-6-nitro-quinoline-3-carbonitrile, 4g (72.35mmol) of iron dust and 8.9g (165.36mmol) of ammonium chloride in 240mL of methanol and water (2:1 ratio) was refluxed for 4hr. The mixture was filtered hot and washed with methanol and water. The product precipitated from the filtrate upon cooling. The solid was collected and dried in vacuo to give 5.8g (79%) yellowish brown solid; ESMS m/z 356.8, 358.8, mp 210-212°C.

Example 46**4-(3-Chloro-4-fluoro-phenylamino)-7-methoxy-6-nitro-quinoline-3-carbonitrile**

A mixture of 4.4 g (16.7 mmol) of 4-chloro-7-methoxy -6-nitro-quinoline-3-carbonitrile and 2.67 g (18.3 mmol) of 3-chloro-4-fluoro aniline in 110 ml of methoxyethanol was refluxed under nitrogen for 4 hours. The reaction mixture was diluted with ethyl acetate and wash with sodium bicarbonate solution and sodium chloride solution. The organic layer was dried over sodium sulfate and then the solvent was removed under vacuum. The residue was chromatographed on silica gel eluting with mixture of ethyl acetate and methanol to give 3 g yellow solid: mass spectrum (electrospray, m/e): 372.9.

- 83 -

Example 47**6-Amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile**

A mixture of 4.88 g (13 mmol) of 4-[(3-chloro-4-fluorophenyl)amino]-7-methoxy-6-nitro-quinoline-3-carbonitrile, 5.2 g (97.5 mmol) of ammonium chloride, and 3.3 g (58.5 mmol) iron was stirred at reflux in 60 ml of water and 60 ml of methanol for 4.5 hours. The mixture was diluted with 500 ml of hot ethyl acetate and the hot mixture was filtered. The filtration was washed with saturated sodium chloride solution and then the organic layer was dried over sodium sulfate. The solvent was removed and the residue was chromatographed on silica gel eluting with mixture of ethyl acetate and methanol to give 3.38 g of yellow solid: mass spectrum (electrospray, m/e): M+H 343.4.

Example 48**4-(3-Bromo-4-fluoro-phenylamino)-7-methoxy-6-nitro-quinoline-3-carbonitrile**

A mixture of 3.52 g (9.7 mmol) of 4-chloro-7-methoxy -6-nitro-quinoline-3-carbonitrile and 2.0 g (10.7 mmol) of 3-bromo-4-fluoro aniline in 150 ml of methoxyethanol was refluxed under nitrogen for 5.5 hours. The reaction mixture was diluted with ethyl acetate and wash with sodium bicarbonate solution and sodium chloride solution. The organic layer was dried with sodium sulfate and then solvent was removed under vacuum. The residue was chromatographed on silica gel eluting with mixture of ethyl acetate and hexane to give the title compound.

Example 49**6-Amino-4-(3-bromo-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile**

A mixture of 2.9 g (6.95 mmol) of 4-[(3-bromo-4-fluorophenyl)amino]-7-methoxy-6-nitro-quinoline-3-carbonitrile, 6.5 g (121.6 mmol) of ammonium chloride and 4.05 g (73 mmol) of iron in 50 ml of water and 50 ml of methanol for 6 hours. The mixture was diluted with hot ethyl acetate and the hot mixture was filtered. The filtration was washed with saturated sodium chloride solution then the organic layer was dried over sodium sulfate. The solvent was removed and the residue was chromatographed on

5

- 84 -

10

silica gel eluting with mixture of ethyl acetate and methanol to give 2.11 g of light yellow solid: mass spectrum (electrospray, m/c): M+H 386.7 and 388.8.

Example 50

15

5 7-Ethoxy-4-hydroxy-quinoline-3-carbonitrile

20

A mixture of 10 g (73 mmol) of 3-ethoxy aniline and 12.3 g (73 mmol) of ethyl (ethoxymethylene) cyanoacetate was heated in 90 ml of Dowther at 140°C for 7 hours. To this mixture was added 250 ml of Dowther. The solution was stirred and refluxed under nitrogen for 12 hours with periodically distilling out the eliminated ethanol. The mixture was cooled to room temperature and the solid was collected and washed with hexane. The crude solid was treated with boiling ethanol and then filtered to give 9.86 g of brown solid: mass spectrum (electrospray, m/c): M+H 214.7.

25

Example 51

30

15 7-Ethoxy-4-hydroxy-6-nitro-quinoline-3-carbonitrile

35

To a suspension of 5 g (23 mmol) of 7-Ethoxy-4-hydroxy-quinoline-3-carbonitrile in 75 ml of trifluoroacetic anhydride was added 5.5 g (69 mmol) of ammonium nitrate over a period of 6 hours at room temperature. Excess anhydride was removed at reduced pressure at 45°C. The residue was stirred with 300 ml of water. The solid was collected and treated with boiling ethanol to give 3.68 g of tan solid: mass spectrum (electrospray, m/e) M+H 259.8.

20

40

Example 52

45

4-Chloro-7-ethoxy-6-nitro-quinoline-3-carbonitrile

50

25 A mixture of 3.45 g (13 mmol) of 7-Ethoxy-4-hydroxy-6-nitro-quinoline-3-carbonitrile, 5.55 g (26 mmol) of phosphorous pentachloride, and 10 ml of phosphorous oxychloride was refluxed for 3 hours. The mixture was diluted with hexane and the solid was collected. The solid was dissolved in 500 ml of ethyl acetate and washed with cold diluted sodium hydroxide solution. The solution was dried over magnesium sulfate and filtered through a pad of silica gel. The solvent

55

5

- 85 -

10

was removed giving 2.1 g of beige solid: mass spectrum (electrospray, m/e) M+H 277.7.

Example 53

15

4-(3-Bromo-phenylamino)-7-ethoxy-6-nitro-quinoline-3-carbonitrile

20

A mixture of 2.1 g (7.6 mmol) of 4-chloro-7-ethoxy-6-nitro-quinoline-3-carbonitrile and 0.91 ml (8.3 mmol) of 3-bromo aniline in 100 ml ethanol was refluxed under nitrogen for 4.5 hours. The reaction mixture was poured into diluted sodium bicarbonate solution. Ethanol was removed under vacuum. The mixture was diluted with ethyl acetate and the organic layer was separated and dried over sodium sulfate. The solution was concentrated and solid was collected and then washed with hexane. Upon drying, 2.6 g of yellow solid obtained: mass spectrum (electrospray, m/e) M+H 412.8 and 414.9.

25

15

Example 54

30

6-Amino-4-(3-bromo-phenylamino)-7-ethoxy-quinoline-3-carbonitrile

35

A mixture of 2.5 g (6 mmol) of 4-[(3-bromophenyl)amino]-7-ethoxy-6-nitro-quinoline-3-carbonitrile, 2.4 g (45 mmol) of ammonium chloride, and 1.5 g (27 mmol) iron was stirred at reflux in 40 ml of water and 40 ml of methanol for 4 hours. The mixture was diluted with 500 ml of hot ethyl acetate and the hot mixture was filtered. The filtration was washed with saturated sodium chloride solution and then the organic layer was dried over sodium sulfate. The solution was concentrated and 1.5 of beige solid was collected: mass spectrum (electrospray, m/e): M+H 382.8 and 384.8.

40

25

Example 55

45

8-Methoxy-4-hydroxy-6-nitro-quinoline-3-carbonitrile

50

A mixture of 12.6 g (75 mmol) of 2-methoxy-4-nitro aniline and 12.7 g (75 mmol) of ethyl (ethoxymethylene) cyanoacetate was heated in 100 ml of Dowther at 120°C for overnight and 180°C for 20 hours. To this mixture was added 300 ml of Dowther. The solution was stirred and refluxed under nitrogen for 12 hours with periodically

55

5

- 86 -

10

distilling out the eliminated ethanol. The mixture was cooled to room temperature and the solid was collected and washed with hexane. The crude solid was treated with boiling ethanol and then filtered to give 12 g of brown solid: mass spectrum (electrospray, m/e): M+H 245.8.

15

5

Example 56

4-Chloro-8-methoxy-6-nitro-quinoline-3-carbonitrile

20

A mixture of 4 g (16 mmol) of 8-Methoxy-4-hydroxy-6-nitro-quinoline-3-carbonitrile, 6.66 g (32 mmol) of phosphorous pentachloride, and 15 ml of phosphorous oxychloride was refluxed for 2.5 hours. The mixture was diluted with hexane and the solid was collected. The solid was dissolved in 500 ml of ethyl acetate and washed with cold diluted sodium hydroxide solution. The solution was dried over magnesium sulfate and filtered through a pad of silica gel. The solvent was removed giving 2.05 g of tan solid: mass spectrum (electrospray, m/e) M+H 263.7.

25

15

30

Example 57

6-nitro-4-(3-bromo-phenylamino)-8-methoxy-quinoline-3-carbonitrile

35

A mixture of 1.9 g (7.6 mmol) of 4-chloro-8-methoxy-6-nitro-quinoline-3-carbonitrile and 0.86 ml (8.3 mmol) of 3-bromo aniline in 95 ml ethanol was refluxed under nitrogen for 5 hours. The reaction mixture was poured into diluted sodium bicarbonate solution. Ethanol was removed under vacuum. The mixture was diluted with ethyl acetate and the organic layer was separated and dried over sodium chloride. The solution was concentrated and solid was collected and then washed with hexane. Upon drying, 2.3 g of yellow solid obtained: mass spectrum (electrospray, m/e) M+H 398.8 and 400.8.

40

25

45

Example 58

6-Amino-4-(3-bromo-phenylamino)-8-methoxy-quinoline-3-carbonitrile

50

A mixture of 2.15 g (5 mmol) of 4-[(3-bromophenyl)amino]-8-methoxy-6-nitro-quinoline-3-carbonitrile, 1.95 g (37.5 mmol) of ammonium chloride, and 1.26 g (22.5

55

5

- 87 -

10

15

mmol) iron was stirred at reflux in 40 ml of water and 40 ml of methanol for 3 hours. The mixture was diluted with 500 ml of hot ethyl acetate and the hot mixture was filtered. The filtration was washed with saturated sodium chloride solution and then the organic layer was dried over sodium sulfate. The solution was concentrated and 0.43 of dark yellow solid was collected: mass spectrum (electrospray, m/e): M+H 368.9 and 370.9.

Example 59

20

4-Chloro-but-2-ynoic acid

25

30

Propargyl chloride (2 mL, 26.84mmol) was dissolved in 40 mL of tetrahydrofuran under nitrogen and cooled to -78°C. After addition of n-butyllithium (5.4 mL, 13.42mmol, 2.5 M in n-hexane) and stirred for 15 min, a stream of dry carbon dioxide was passed through it at -78°C for two hours. The reaction solution was filtered and neutralized with 3.5 mL of 10% sulfuric acid. After evaporation of the solution, the residue was extracted with ether. The ether solution was washed with saturated brine solution, and dried over sodium sulfate. After evaporation of the dry ether solution, 0.957g (60%) of an oil product was obtained: ESMS m/z 116.6 (M-H⁺).

35

20

Example 60

4-Dimethylamino -but-2-ynoic acid

40

45

50

30

n-Butyl lithium in hexane (96mL, 2.5 M in n-hexane) was slowly added to 1-dimethylamino-2-propyne (20g, 240mmol) in 100 mL of tetrahydrofuran under nitrogen. The mixture was stirred for 1 h at -78°C, then dry carbon dioxide was pass through overnight. The resulting solution was poured into water and washed with ethyl acetate. The aqueous layer was evaporated under reduced pressure to give the crude acid. The dry acid was dissolved in methanol, and the insoluble salt was removed via filtration. The filtrate was collected and dried in vacuo to give 15.6g of 4-dimethylamino -but-2-ynoic acid: mass spectrum (m/e):M-H 126.

55

5

- 88 -

Example 61

10

Bis-(2-methoxy-ethyl)-prop-2-ynyl-amine

15

20

10

Propargyl bromide (17.8g, 150mmol) was added dropwise to a mixture of bis(2-methoxy-ethyl)amine (20g, 150mmol) and cesium carbonate (49g, 150mmol) in 350mL of acetone. The mixture was stirred overnight under nitrogen at room temperature. The inorganic salts were then filtered off, and the solvent was removed. The residue was dissolved in saturated sodium bicarbonate solution and extracted with ethyl acetate. The organic extracts were then evaporated to give 20g of bis-(2-methoxy-ethyl)-prop-2-ynyl-amine: mass spectrum (m/e): M+H 172.

Example 62

25

4-[Bis-(2-methoxy-ethyl)-amino]-but-2-ynoic acid

30

35

20

n-Butyl lithium in hexane (42mL, 2.5M in n-hexane) was slowly added to bis-(2-methoxy-ethyl)-prop-2-ynyl-amine (18g, 105mmol) in 80mL of tetrahydrofuran under nitrogen. The mixture was stirred for 1 h at -78°C, then dry carbon dioxide was passed through overnight. The resulting solution was poured into water and washed with ethyl acetate. The aqueous layer was evaporated under reduced pressure to give the crude acid. The dry acid was dissolved in methanol, and the insoluble salt was removed via filtration. The filtrate was collected and dried in vacuo to give 18g of 4-[bis-(2-methoxy-ethyl)-amino]-but-2-ynoic acid: mass spectrum (m/e):M-H 214.

Example 63

40

1-Methyl-4-prop-2-ynyl-piperazine

45

50

30

Propargyl bromide (23.8g, 200mmol) was added dropwise to a mixture of 1-methyl-piperazine (20g, 200mmol) and cesium carbonate (65g, 200mmol) in 350mL of acetone. The mixture was stirred overnight under nitrogen at room temperature. The inorganic salts were then filtered off, and the solvent was removed. The residue was dissolved in saturated sodium bicarbonate solution and extracted with ethyl acetate. The organic extracts were then evaporated to give 7.5g of 1-methyl-4-prop-2-ynyl-piperazine: mass spectrum (m/e): M+H 139.

55

- 89 -

Example 644-(4-Methyl-piperazin-1-yl)-but-2-ynoic acid

n-Butyl lithium in hexane (17.2mL, 2.5M in n-hexane) was slowly added to 1-methyl-4-prop-2-ynyl-piperazine (6.0g, 43.5mmol) in 40mL of tetrahydrofuran under nitrogen. The mixture was stirred for 1 hr at -78°C, then dry carbon dioxide was passed through overnight. The resulting solution was poured into water and washed with ethyl acetate. The aqueous layer was evaporated under reduced pressure to give the crude acid. The dry acid was dissolved in methanol, and the insoluble salt was removed via filtration. The filtrate was collected and dried in vacuo to give 7g of 4-(4-methyl-piperazin-1-yl)-but-2-ynoic acid: mass spectrum (m/e): M-H 181.

Example 65(2-Methoxy-ethyl)-methyl-prop-2-ynyl-amine

Propargyl bromide (26.8g, 225mmol) was added dropwise to a mixture of N-(2-methoxyethyl)methyl amine (20g, 225mmol) and cesium carbonate (73g, 225mmol) in 350mL of acetone. The mixture was stirred overnight under nitrogen at room temperature. The inorganic salts were then filtered off, and the solvent was removed. The residue was dissolved in saturated sodium bicarbonate solution and extracted with ethyl acetate. The organic extracts were then evaporated to give 14g of (2-methoxy-ethyl)-methyl-prop-2-ynyl-amine: mass spectrum (m/e): M+H 127.

Example 664-[(2-Methoxy-ethyl)-methyl-aminol]-but-2-ynoic acid

n-Butyl lithium in hexane (37.8mL, 2.5 M in n-hexane) was slowly added to (2-methoxy-ethyl)-methyl-prop-2-ynyl-amine (12.0g, 94.5mmol) in 90mL of tetrahydrofuran under nitrogen. The mixture was stirred for 1 hr at -78°C, then dry carbon dioxide was passed through overnight. The resulting solution was poured into water and washed with ethyl acetate. The aqueous layer was evaporated under reduced pressure to give the crude acid. The dry acid was dissolved in methanol, and the insoluble salt was removed via filtration. The filtrate was collected and dried in

- 90 -

vacuo to give 15g of 4-[(2-methoxy-ethyl)-methyl-amino]-but-2-ynoic acid: mass spectrum (m/e): M-H 170.

Example 67

Allyl-methyl-prop-2-ynyl-amine

Propargyl bromide (33.4g, 281mmol) was added dropwise to a mixture of isopropyl-methyl-amine (20g, 281mmol) and cesium carbonate (90g, 281mmol) in 350mL of acetone. The mixture was stirred overnight under nitrogen at room temperature. The inorganic salts were then filtered off, and the solvent was removed. The residue was dissolved in saturated sodium bicarbonate solution and extracted with ethyl acetate. The organic extracts were then evaporated to give 4.6g of allyl-methyl-prop-2-ynyl-amine: mass spectrum (m/c): M+H 110.

Example 68

4-(Allyl-methyl-amino)-but-2-ynoic acid

n-Butyl lithium in hexane (16.4mL, 2.5M in n-hexane) was slowly added to allyl-methyl-prop-2-ynyl-amine (4.5g, 46mmol) in 50 mL of tetrahydrofuran under nitrogen. The mixture was stirred for 1 hr at -78°C, then dry carbon dioxide was passed through overnight. The resulting solution was poured into water and washed with ethyl acetate. The aqueous layer was evaporated under reduced pressure to give the crude acid. The dry acid was dissolved in methanol, and the insoluble salt was removed via filtration. The filtrate was collected and dried in vacuo to give 4.1g of 4-(allyl-methyl-amino)-but-2-ynoic acid: mass spectrum (m/e): M-H 152.

Example 69

4-Methoxymethoxy-but-2-ynoic acid

To a suspension of 8.2 g of 60% sodium hydride in mineral oil in 271 mL of tetrahydrofuran at 0°C with stirring under nitrogen was added dropwise 10 g of propargyl alcohol over 15 min. The mixture was stirred an additional 30 min. To the stirred mixture at 0°C was added 15.8 g of chloromethylmethyl ether. Stirring was continued at room temperature over night. The mixture was filtered and the solvent

5

- 91 -

10

15

20

was removed from the filtrate. The residuc was distilled (35-38°C, 4 mm) giving 8.5 g of a liquid. The distillate was dissolved in 200 mL of ether. The solution was stirred under nitrogen and cooled to -78°C as 34.1 mL of 2.5 molar n-butyl lithium in hexanes was added over 15 min. Stirring was continued for another 1.5 hr. Dry carbon dioxide was allowed to pass over the surface of the stirring reaction mixture as it warmed from -78°C to room temperature. The mixture was stirred under a carbon dioxide atmosphere over night. The mixture was poured into a mixture of 14 mL of hydrochloric acid and 24 mL of water. The organic layer was separated and dried over magnesium sulfate. The solvent was removed and the residue was maintained at 100°C at 4 mm for 1 hr giving 10.4 g 4-Methoxymethoxy-but-2-ynoic acid.

25

Example 70

4-Bromo crotonic acid

30

35

40

After the method of Braun [Giza Braun, J. Am. Chem. Soc. 52, 3167 (1930)], 11.76 mL (17.9 grams 0.1 moles) of methyl 4-bromo crotonate in 32 mL of ethanol and 93 mL of water was cooled to -11 °C. The reaction was stirred vigorously, and 15.77 g (0.05 moles) of finely powdered barium hydroxide was added portionwise over a period of about an hour. Cooling and vigorous stirring were continued for about 16 hours. The reaction mixture was then extracted with 100 mL of ether. The aqueous layer was treated with 2.67 mL (4.91 g; 0.05 moles) of concentrated sulfuric acid. The resulting mixture was extracted with 3-100 mL portions of ether. The combined ethereal extracts were washed with 50 mL of brine, then dried over sodium sulfate. The solution was taken to an oil *in vacuo*. This oil was taken up in about 400 mL of boiling heptane, leaving a gum. The heptane solution was separated and boiled down to about 50mL. Cooling gave 3.46 g of product.

45

Example 71

4-(2-Methoxy-ethoxy)-but-2-ynoic acid

50

To a suspension of 6.04 g (151 mmol) of 60% sodium hydride in 200 ml of tetrahydrofuran at 0 °C was add 10 g (131.4 mmol) of 2-methoxyethanol dropwise over 15 min. After 1 hr, 19.54 g (131.4 mmol) of 80% propargyl bromide was added

55

5

- 92 -

10

15

20

dropwise. After stirring 17 hr at room temperature, the mixture was filtered and the solvent was removed. The residue was distilled (48-51 °C, 4mm) to give 11.4 g of a colorless liquid. This was dissolved in 250 ml of ether and cooled to -78 °C with stirring under nitrogen. To this solution was added 39.95 ml (99.9 mmol) of 2.5M n-butyl lithium solution in hexanes dropwise over 15 min. After 1.5 hr, dry carbon dioxide was bubbled in as the mixture slowly warmed to room temperature. The mixture was maintained in a carbon dioxide atmosphere overnight. To the mixture was added 100 ml of 3N hydrochloric acid and solid sodium chloride. The organic layer was separated and dried over magnesium sulfate. The solvent was removed and the residue was maintained under vacuum giving 11.4 g of the title compound. : mass spectrum (electrospray, m/e, negative mode): M-H 156.8.

25

Example 72

4-(2-Methoxy-ethoxy)-but-2-ynoic acid [4-(3-bromo-phenylamino)-3- cyano-quinolin-6-yl]-amide

15

30

35

20

40

25

45

To a solution of 0.56 g (3.54 mmol) of 4-(2-methoxy-ethoxy)-but-2-ynoic acid and 0.46 g (3.4 mmol) of isobutyl chloroformate in 12 ml of tetrahydrofuran was added at 0°C with stirring 0.36 g (3.54 mmol) of N-methylmorpholine. After 15 min, 1.0 g (2.95 mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile was added. After stirring 3 hr at 0°C and 17 hr at room temperature, the mixture was poured into a saturated solution of sodium bicarbonate. The mixture was extracted with ethyl acetate and the organic layer was dried over magnesium sulfate. The solvent was removed and the residue was purified by chromatography on silica gel eluting with chloroform- ethyl acetate mixtures to give 0.53 g of 4-(2-Methoxy-ethoxy)-but-2-ynoic acid [4-(3-bromo-phenylamino)-3- cyano-quinolin-6-yl]-amide as a yellow powder: mass spectrum (electrospray, m/e.): M+H 480.9.

Example 73

4-(Methoxymethoxy)-but-2-ynoic acid

50

30

To a suspension of 8.2 g (205 mmol) of 60% sodium hydride in 271 ml of tetrahydrofuran was added dropwise at 0 °C with stirring 10.0 g (178.4 mmol) of

55

5

- 93 -

10

15

20

propargyl alcohol . After 30 min, 15.8 g (196.2 mmol) of chloromethylmethyl ether was added. After stirring over the weekend at room temperature, the mixture was filtered and the solvent was remove. The residue was distilled (35-38 °C, 4mm) to give 8.54 g of a colorless liquid. This was dissolved in 200 ml of ether and cooled to -78 °C with stirring under nitrogen. To this solution was added 34.1 ml (85.3 mmol) of 2.5M n-butyl lithium solution in hexanes dropwise over 15 min. After 1.5 hr, dry carbon dioxide was bubbled in as the mixture slowly warmed to room temperature. The mixture was maintained in a carbon dioxide atmosphere overnight. To the mixture was added 14 ml of hydrochloric acid in 24 ml water. The organic layer was separated and dried over magnesium sulfate. The solvent was removed and the residue was maintained under vacuum giving 10.4 g of the title compound as a liquid.

25

Example 74

4-Methoxymethoxy-but-2-ynoic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide

15

30

35

20

40

25

45

To a solution of 0.51 g (3.54 mmol) of 4-(methoxymethoxy)-but-2-ynoic acid and 0.46 g (3.4 mmol) of isobutyl chloroformate in 12 ml of tetrahydrofuran was added at 0°C with stirring 0.36 g (3.54 mmol) of N-methylmorpholine. After 15 min, 1.0 g (2.95 mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile was added. After stirring 3 hr at room temperature, the mixture was poured into a saturated solution of sodium bicarbonate. The mixture was extracted with ethyl acetate and the organic layer was dried over magnesium sulfate. The solvent was removed and the residue was purified by chromatography on silica gel eluting with chloroform- ethyl acetate mixtures to give 0.66g of 4-methoxymethoxy-but-2-ynoic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide as a yellow powder: mass of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile :mass spectrum (electrospray, m/e,): M+H 465.1, 467.0.

50

55

- 94 -

Example 75

N-[4-[(3-Bromophenyl)amino]-3-cyano-6-quinolinyl]-4-(bis-(2-methoxyethyl)amino)-2-butyramide

Isobutyl chloroformate (0.785g, 5.75mmol) was dropwise added into an ice cold solution of 4-(bis-methoxyethylamino)-2-butyric acid (1.9g, 8.85mmol) and N-methylmorpholine (0.9386g, 9.28mmol) in 50mL of tetrahydrofuran under N₂. After stirring for 30min, a solution of 1.5g (4.42mmol) of 6-amino-4-[(3-bromophenyl)-amino]-quinoline-3-carbonitrile in 10mL of pyridine was added dropwise and the mixture was stirred at 0°C for 2hr. The reaction was quenched with ice water, poured into saturated sodium bicarbonate and brine, and extracted with ethyl acetate. The ethyl acetate layer was concentrated and purified by flash column chromatography. The product fractions were collected, and dried in vacuo to give 0.82 (35%) of light brown solid; ESMS m/z 536.1, 538.1 (M+H⁺); mp 98-101°C.

Example 76

N-[4-[(3-Bromophenyl)amino]-3-cyano-6-quinolinyl]-4-(N-methoxyethyl-N-methylamino)-2-butyramide

Isobutyl chloroformate (0.785g, 5.75mmol) was dropwise added into an ice cold solution of 4-(N-methoxyethyl-N-methylamino)-2-butyric acid (1.5g, 8.84mmol) and N-methylmorpholine (1.36g, 13.3mmol) in 60mL of tetrahydrofuran under N₂. After stirring for 30min, a solution of 1.5g (4.42mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile in 15mL of pyridine was added dropwise and the mixture was stirred at 0°C for 2hr. The reaction was quenched with ice water, poured into saturated sodium bicarbonate and brine, and extracted with ethyl acetate. The ethyl acetate layer was concentrated and purified by flash column chromatography. The product fractions were collected, and dried in vacuo to give 0.32 (15%) of reddish brown solid; ESMS m/z 492.0, 494.0 (M+H⁺); mp 95°C (dec).

- 95 -

Example 77**N-[4-[(3-Bromophenyl)amino]-3-cyano-6-quinolinyl]-4-(N-allyl-N-methylamino)-2-butynamide**

Isobutyl chloroformate (0.785g, 5.75mmol) was dropwise added into an ice cold solution of 4-(N-allyl-N-methylamino)-2-butynoic acid (1.4g, 8.84mmol) and N-methylmorpholine (0.94g, 9.3mmol) in 80mL of tetrahydrofuran under N₂. After stirring for 30min, a solution of 1.5g (4.42mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile in 15mL of pyridine was added dropwise and the mixture was stirred at 0°C for 2hr. The reaction was quenched with ice water, poured into saturated sodium bicarbonate and brine, and extracted with ethyl acetate. The ethyl acetate layer was concentrated and purified by flash column chromatography. The product fractions were collected, and dried in vacuo to give 0.60 (29%) of brown solid; ESMS m/z 474.4, 476.4 (M+H⁺); mp 133-135°C.

Example 78**1-Methyl-1,2,5,6-tetrahydro-pyridine-3-carboxylic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide**

To a solution of 1 g (2.95 mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile and 1.9 g (14.7 mmol) of diisopropylethylamine was stirred in 19 ml of tetrahydrofuran and solid N-methyl-1,2,5,6-tetrahydronicotinyl chloride hydrochloride was added portionwise at 0°C. After stirring 1 hr at 0°C and 2 hr at room temperature, the mixture was poured into saturated sodium bicarbonate and extracted with ethylacetate. The solution was dried over magnesium sulfate. Solvent was removed and the residue was recrystallized from methanol-ethylacetate giving 0.92 g of a yellow powder : mass spectrum (electrospray, m/e,): M+H 462.4, 464.4.

Example 79**4-((2S)-2-Methoxymethylpyrrolidin-1-yl)but-2-ynoic Acid [4-(3-bromophenylamino)-3-cyanoquinolin-6-yl]amide**

To an ice cold solution of 1.46 g (7.40 mmol) of 4-((2S)-2-methoxymethylpyrrolidin-1-yl)but-2-ynoic acid in 85 mL of THF under N₂ was added 0.897 g (8.88 mmol) of

5

- 96 -

10

15

20

N-methylmorpholine and 0.811 g (5.92 mmol) of isobutyl chloroformate. After stirring in the cold for 30 min, a solution of 1.00 g (2.96 mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile in 8 mL of pyridine was added dropwise. The reaction was slowly warmed to 25°C over 3 h. The reaction was poured into ice water, satd NaHCO₃ was added and the product was extracted with ethyl acetate. After drying and solvent evaporation, the residue was chromatographed on silica gel (10% methanol in ethyl acetate). The yield was 0.560 g of 4-((2S)-2-methoxymethylpyrrolidinyl-1-yl)but-2-ynoic acid [4-(3-bromophenylamino)-3-cyanoquinolin-6-yl]amide as a brown foam: mass spectrum (electrospray, m/e): M+H 518.0, 520.0.

25

Example 80

4-((2S)-2-methoxymethylpyrrolidin-1-yl)butynoic Acid

30

35

n-Butyllithium solution in hexane (35.9 mmol) was added over 10 min to a solution of 5.49 g (35.9 mmol) of (2S)-2-methoxymethyl-1-prop-2-ynylpyrrolidine in 100 mL of THF at -78°C under N₂. After stirring cold for 1 h, CO₂ was bubbled into the solution as it slowly came to 25°C. After stirring overnight, 100 mL of water was added, the reaction was extracted with ethyl acetate and the extracts were discarded. The reaction was adjusted to pH 7 with 20% H₂SO₄ and solvent was removed. The residue was slurried with methanol and filtered. The filtrate was evaporated and dried in vacuo to give 7.06 g of 4-((2S)-2-methoxymethylpyrrolidin-1-yl)butynoic acid as a brown foam: mass spectrum (electrospray, m/e): M+H 198.0.

40

Example 81

25

(2S)-2-Methoxymethyl-1-prop-2-ynylpyrrolidine

45

50

A mixture of 4.82 g (41.9 mmol) of S-2-(methoxymethyl)pyrrolidine, 13.7 g (41.9 mmol) of cesium carbonate and 5.00 g (41.9 mmol) of propargyl bromide in 80 mL of acetone was stirred at 25°C overnight. The reaction was filtered and solvent was removed from the filtrate. The residue was diluted with a small amount of water and satd NaHCO₃ and extracted with ether. The extract was treated with Darco, dried and

55

- 97 -

evaporated to give 5.93 g of (2S)-2-methoxymethyl-1-prop-2-ynylpyrrolidine as a yellow orange oil: mass spectrum (electrospray, m/e): 153.8.

Example 82

4-(1,4-Dioxo-8-azaspiro[4,5]dec-8-yl)but-2-ynoic Acid [4-(3-Bromophenylamino)-3-cyanoquinolin-6-yl] Amide

To an ice cold solution of 1.75 g (7.78 mmol) of 4-(1,4-dioxo-8-azaspiro[4,5]dec-8-yl)but-2-ynoic acid in 100 mL of THF under N₂ was added 0.942 g (9.33 mmol) of N-methylmorpholine followed by 0.852 g (6.22 mmol) of isobutyl chloroformate. After stirring in the cold for 30 min, a solution of 1.05 g (3.11 mmol) of 6-amino-4-[(3-bromophenyl)amino]-quinoline-3-carbonitrile in 8 mL of pyridine was added dropwise. After stirring in the cold for 5 h, the reaction was poured into ice water and satd NaHCO₃ was added. The mixture was extracted with ethyl acetate and the extracts were dried and evaporated. Chromatography of the residue on silica gel (20% methanol in ethyl acetate) gave 0.590 g of 4-(1,4-dioxo-8-azaspiro[4,5]dec-8-yl)but-2-ynoic acid [4-(3-bromophenylamino)-3-cyanoquinol-6-yl] amide as a brown foam: mass spectrum (electrospray, m/e): M+H 546.0, 548.1.

Example 83

4-(1,4-Dioxo-8-azaspiro[4,5]dec-8-yl)but-2-ynoic Acid

n-Butyllithium in hexane (55.8 mmol) was added dropwise to a solution of 10.1 g (55.8 mmol) of 3-(1,4-dioxo-8-azaspiro[4,5]dec-8-yl)but-2-yne in 185 mL of THF at -78°C under N₂. After stirring at -78°C for 1 h, CO₂ was bubbled into the solution as it slowly came to 25°C. After stirring overnight, the reaction was diluted with 150 mL of water, extracted with ethyl acetate and the extracts were discarded. The solution was adjusted to pH 6 with 2 M sulfuric acid and evaporated. The residue was slurried with methanol and filtered. The filtrate was evaporated and dried in vacuo to give 4.5 g of 4-(1,4-dioxo-8-azaspiro[4,5]dec-8-yl)but-2-ynoic acid as a brown amorphous solid: mass spectrum electrospray, m/e): M+H 225.8.

- 98 -

Example 843-(1,4-Dioxo-8-azaspiro[4,5]dec-8-yl)but-2-yne

A mixture of 10.0 g (69.9 mmol) of 1,4-dioxo-8-azaspiro[4,5]decane, 22.8 g (69.9 mmol) of cesium carbonate and 8.32 g (69.9 mmol) of propargyl bromide in 165 mL of acetone was stirred overnight at 25 °C. The reaction was filtered and the filtrate was evaporated to dryness. A small amount of water and satd NaHCO₃ was added to the residue and it was extracted with ether. The ethereal extracts were treated with Darco, dried and evaporated to give 10.8 g of 3-(1,4-dioxo-8-azaspiro[4,5]dec-8-yl)but-2-yne as a yellow orange oil: mass spectrum (electrospray, m/e): M+H 181.8.

Example 854-(3-Bromo-phenylamino)-6-(2-ethoxy-3,4-dioxo-cyclobut-1-enylamino)-quinoline-3-carbonitrile

A mixture of 1.00 g (2.95 mmol) 6-amino-4-(3-chloro-phenylamino)-quinoline-3-carbonitrile, 20 mL ethanol, and 0.873 mL (5.90 mmol) 3,4-diethoxy-3-cyclobutene-1,2-dione was heated to reflux under N₂. At 4 hours removed heat, cooled to 25°C and stirred overnight. Decanted off solution and stripped solvent. Added ether to crystallize, collected solids and dried. Boiled in ethyl acetate to remove cyclobutene starting material. Dried in vacuo, giving 249 mg of yellow solid: mass spectrum (electrospray m/e): M+H = 463.2.

Example 864-(4-Chloro-2-fluoro-phenylamino)-6,7-dimethoxy-quinoline-3-carbonitrile

A mixture of 2.0 g of 4-chloro-6,7-dimethoxy-quinoline-3-carbonitrile, 1.46 g of 4-chloro-2-fluoroaniline, 0.925 g of pyridine hydrochloride, and 125 ml of ethoxyethanol was stirred under nitrogen, at reflux temperature for 1 h. The mixture was cooled and added to 1000 ml of water. To this mixture was added sodium carbonate to pH 9. The product was collected, washed with water, and dried to give 2.61 g of 4-(4-chloro-2-fluoro-phenylamino)-6,7-dimethoxy-quinoline-3-carbonitrile as a solid, mp 139-141°C; mass spectrum (electrospray, m/e): M+H 357.9.

- 99 -

Example 874-(4-Chloro-2-fluoro-phenylamino)-6,7-dihydroxy-quinoline-3-carbonitrile

A mixture of 0.358 g of 4-(4-chloro-2-fluoro-phenylamino)-6,7-dimethoxy-quinoline-3-carbonitrile and 3 g of pyridine hydrochloride was stirred under nitrogen at 210 -220°C for 20 minutes. The mixture was cooled and added to 50 ml of 3% ammonium hydroxide solution. The product was collected, washed with water, and dried to give 0.302 g of 4-(4-chloro-2-fluoro-phenylamino)-6,7-dihydroxy-quinoline-3-carbonitrile as a solid, mp 270-272°C; mass spectrum (EI, m/e): M 329.0363.

Example 884-(4-Chloro-2-fluoro-phenylamino)-6-methoxy-7-(2-pyridin-4-yl-ethoxy)-quinoline-3-carbonitrile

To a solution of 0.655 g of triphenylphosphine in 20 ml of tetrahydrofuran was added dropwise 0.348 mg of diethyl azodicarboxylate. The solution was stirred for one minute and added to a mixture of 0.330 g of 4-(4-chloro-2-fluoro-phenylamino)-6,7-dihydroxy-quinoline-3-carbonitrile and 0.500 g of 2-(4-pyridyl)ethanol in 100 ml of tetrahydrofuran. The mixture was stirred at room temperature for 4 hours, and 10 ml of methanol was added.

To a solution of 0.655 g of triphenylphosphine in 20 ml of tetrahydrofuran was added dropwise 0.348 mg of diethyl azodicarboxylate. The solution was stirred for one minute and added to the above mixture. The mixture was stirred overnight and concentrated in vacuo. The residue was chromatographed on silica gel eluting with 5% methanol in dichloromethane. Solvent was removed from product fractions giving 0.034 g of 4-(4-Chloro-2-fluoro-phenylamino)-6-methoxy-7-(2-pyridin-4-yl-ethoxy)-quinoline-3-carbonitrile as a white wax: mass spectrum (EI, m/e): M 448.1104. The regiochemistry was assigned unequivocally by NMR analysis (Proton, DQF-COSY, NOESY, {¹H-¹³C}-HMQC, {¹H-¹³C}-HMBC, {¹H-¹⁵N}-HMBC.

- 100 -

Example 89

4-[2-Methoxy-ethyl)-methyl-aminol-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide dihydrochloride

To a stirring solution of 1.2 g (3.5 mmol) of 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile and 0.52 g (4.0 mmol) of diisopropylethylamine in 40 ml of tetrahydrofuran at 0°C was added a solution of 4-bromo crotonyl chloride in 10 ml of tetrahydrofuran. After 45 min, 1.87 (21 mmol) of 2-methoxyethyl methyl amine was added. After 1hr at room temperature, the mixture was poured into a solution of sodium bicarbonate and extracted with ethyl acetate. The organic solution was dried over magnesium sulfate. The solvent was removed and the residue was purified by chromatography on silica gel. Product eluted with ethyl acetate-methanol-triethylamine 40:8:1 giving 0.87 g of the free base. This was dissolved in 20 ml of ethyl acetate and 10 ml of a solution of hydrogen chloride in ether was added. The solid was collected giving 1.02 g of the title compound a yellow powder: mass spectrum (electrospray, m/e): M+H 498.0, (M+2H)⁺ 248.5.

Example 90

(S)-4-(2-Methoxymethyl-pyrrolidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide dihydrochloride

By using the method of Example 89, 1.2 g (3.5 mmol) of 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile and 2.4 g (21 mmol) of (S)-(+)-2-(methoxymethyl) pyrrolidine were converted to 1.5 g the title compound, obtained as a yellow powder: mass spectrum (electrospray, m/e): M+H 524.0, (M+2H)⁺ 262.4. This reaction can also be done with (R)-(-)-2-(methoxymethyl) pyrrolidine or racemic 2-(methoxymethyl) pyrrolidine giving the R-entaniomer or the racemate, respectively.

5

- 101 -

Example 91

10

4-(3-Hydroxymethyl-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide hydrochloride

15

By using the method of Example 89, 1.1 g (3.2 mmol) of 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile and 2.2 g (19.2 mmol) of 3-hydroxymethyl-piperidine were converted to 0.76 g the title compound, obtained as a yellow powder: mass spectrum (electrospray, m/e): M+H 524.0, (M+2H)⁺ 262.3.

20

Example 92

10

4-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

25

By using the method of Example 89, 1.05 g (3.06 mmol) of 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile and 2.6 g (18.4 mmol) of 1,4-dioxo-8-azaspiro[4.5]decane were converted to 0.62 g the title compound. The free base obtained as a yellow foam: mass spectrum (electrospray, m/e): M+H 552.0, (M+2H)⁺ 270.5.

30

Example 93

35

20

4-(2-Hydroxymethyl-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

40

25

By using the method of Example 89, 1.05 g (3.06 mmol) of 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile and 2.1 g (18.4 mmol) of 2-hydroxymethyl-piperidine were converted to 0.67 g the title compound. The free base was obtained as an off-white powder: mass spectrum (electrospray, m/e): M+H 524.3, (M+2H)⁺ 267.7.

45

Example 94

50

30

4-Bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide.

By using the method of Example 15, 4-bromo crotonyl chloride and 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-quinoline-3-carbonitrile was converted to

55

- 102 -

the title compound which was obtained as a solid that could be purified with boiling methanol.

Example 95

3-{3-[4-(3-Chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]carbonyl-allyl}-5-methyl-thiazol-3-ium bromide

A solution of 0.5 g (1 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and 0.6 g (6.1 mmol) of 5-methyl thiazole was refluxed for 4 hr. The mixture was diluted with ethyl acetate and cooled. Solid was collected and recrystallized from methanol-acetone-ethyl acetate to give 0.2 g of the title compound as a yellow powder: mass spectrum (electrospray, m/e): M⁺ 508.0, 509.9, (M+H)⁺ 254.4, 255.1.

Example 96

3-{3-[4-(3-Chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]carbonyl-allyl}-4-methyl-thiazol-3-ium bromide

A solution of 0.7 g (1.4 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and 0.85 g (8.6 mmol) of 4-methyl thiazole was refluxed for 17 hr. The mixture was diluted with ethyl acetate and cooled. Solid was collected and recrystallized from methanol-acetone-ethyl acetate to give 0.3 g of the title compound as a yellow powder: mass spectrum (electrospray, m/e): M⁺ 507.9, 509.8, (M+H)⁺ 254.4, 255.1.

Example 97

Methyl 4-benzyloxy-2-(dimethylaminomethyleneamino)-5-methoxybenzoate

A stirred mixture of 70.0g (244 mmol) of methyl 2-amino-4-benzyloxy-5-methoxybenzoate (Phytochemistry **1976**, 15, 1095) and 52 ml of dimethylformamide dimethyl acetal was heated at 100°C for 1.5 h, cooled, and evaporated directly under high vacuum to give 81.3 g of off-white solid, mp 134-140°C; NMR (CDCl₃) δ 3.01 (s, Me₂N).

- 103 -

Example 987-benzyloxy-4-hydroxy-6-methoxy-quinoline-3-carbonitrile

To a stirred solution of 26.9 ml of *n*-butyllithium (2.5 M in hexane) in 50 ml of THF at -78°C was added a 3.51 ml of acetonitrile in 20 ml of THF during 10 min. After stirring at -78°C for 30 min, the mixture was treated with 10 g of methyl 4-benzyloxy-2-(dimethylaminomethyl)camino-5-methoxybenzoate in 20 ml of THF during 5 min. After 15 min at -78°C the stirred mixture was warmed to 0°C for a further 30 min. It was then treated with 5 ml of acetic acid, warmed to 25°C and stirred for 30 min. The mixture was evaporated to dryness, and diluted with aqueous sodium bicarbonate. The resulting off-white solid was filtered, washed with water, ethyl acetate and ether. After drying, 4.5 g of 7-benzyloxy-4-hydroxy-6-methoxy-quinoline-3-carbonitrile was obtained as an off-white solid, dec > 255°C ; mass spectrum (electrospray, m/e) M+H 307.

15

Example 997-benzyloxy-4-chloro-6-methoxy-quinoline-3-carbonitrile

To a stirred suspension of 1 g of 7-benzyloxy-4-hydroxy-6-methoxy-quinoline-3-carbonitrile in 10 ml of methylene chloride was added 5 ml of oxalyl chloride (2M in methylene chloride), and 2 drops of N,N-dimethylformamide. The mixture was refluxed for 20 min and to it was slowly added aqueous sodium bicarbonate until the bubbling ceased. Following separation of the layers, the organic layer was evaporated to a small volume, then passed through a plug of magneson. Elution with 50 ml methylene chloride, followed by evaporation provided 0.6 g of 7-benzyloxy-4-chloro-6-methoxy-quinoline-3-carbonitrile as a pale yellow solid, mp 282-284°C; mass spectrum (electrospray, m/e) M+H 325.

45

Example 1004-chloro-7-hydroxy-6-methoxy-quinoline-3-carbonitrile

A stirred suspension of 0.54 g of 7-benzyloxy-4-chloro-6-methoxy-quinoline-3-carbonitrile in 10 ml of methylene chloride was cooled to 0°C. To this was added 10 ml of boron trichloride (1M in methylene chloride). The mixture darkened as it

55

5

- 104 -

10

15

20

10

warmed to room temperature and a solid precipitated out. After stirring for 1 hour, no further reaction was observed. The solid (unreacted starting material) was filtered off, the remaining solution was cooled to 0°C and quenched by the dropwise addition of methanol. Following evaporation of the solvent, the residue was dissolved in methylene chloride/methanol/acetone. Purification of this residue was carried out using silica gel chromatography, eluting with a solvent gradient of 1 to 5 percent methanol/methylene chloride, to provide 0.075 g of 4-chloro-7-hydroxy-6-methoxy-quinoline-3-carbonitrile as a yellow solid, dec >245°C; mass spectrum (electrospray, m/e) M+H 235.2.

Example 101

25

4-chloro-6-methoxy-7-(3-pyridin-4-yl-propoxy)-quinoline-3-carbonitrile

30

35

20

A mixture of 0.070 g of 4-chloro-7-hydroxy-6-methoxy-quinoline-3-carbonitrile, 0.062 g of 3-(4-pyridyl)-1-propanol and 0.235 g of triphenylphosphine in 3 ml of methylene chloride under nitrogen was cooled to 0°C. To this was added 0.14 ml of diethyl azodicarboxylate dropwise. After 30 minutes, the reaction mixture was warmed to room temperature and further stirred for 2 hours. The mixture was concentrated down to 1ml and purified by silica gel chromatography, eluting with a solvent gradient of 1 to 2 percent methanol/methylene chloride, to provide 0.090 g of 4-chloro-6-methoxy-7-(3-pyridin-4-yl-propoxy)-quinoline-3-carbonitrile as an off-white gum.

40

Example 102

25

4-(3-hydroxy-4-methyl-phenylamino)-6-methoxy-7-(3-pyridin-4-yl-propoxy)-quinoline-3-carbonitrile

45

50

30

A mixture of 0.090 g of 4-chloro-6-methoxy-7-(3-pyridin-4-yl-propoxy)-quinoline-3-carbonitrile, 0.050 g of 3-hydroxy-4-methylaniline, 0.039 g of pyridine hydrochloride and 3 ml of ethoxyethanol was stirred under nitrogen at reflux temperature for 20 minutes. The mixture was cooled and filtered. The product was washed with saturated sodium bicarbonate, water, then dried to give 0.080 g of 4-(3-hydroxy-4-methyl-phenylamino)-6-methoxy-7-(3-pyridin-4-yl-propoxy)-quinoline-3-carbonitrile

55

- 105 -

hydrochloride as a solid, dec >153°C ; mass spectrum (electrospray, m/e): M+H 440.9.

Example 103

4-Diallylamino-but-2-enoic acid

[4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

A solution of 0.24 g (0.5mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide in 1 ml of N,N-dimethylformamide and 4 ml of tetrahydrofuran was stirred with 0.49 ml (4 mmol) of diallylamine for 3 hr. The reaction was quenched with 10 ml of saturated sodium bicarbonate and 10 ml of ethyl acetate. The insoluble precipitate was collected and washed with water to give 23.7 mg of the title compound (free base); mass spectrum (electrospray, m/e): M+H 506.0. The ethyl acetate layer was washed with water. After the solvent was removed, the crude product was purified by preparative HPLC (C18 column, Gradient from 2% acetonitrile containing 0.05% trifluoroacetic acid to 100% acetonitrile containing 0.05% trifluoroacetic acid in 12 min) to yield 97.9 mg of the product as the bis-trifluoroacetate salt ; mass spectrum (electrospray, m/e): M+H 506.0.

Example 104

4-[Bis-(2-methoxy-ethyl)-amino]-but-2-enoic acid

[4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

In the manner of Example 103, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and bis-(2-methoxyethyl)-amine was converted to 52.3 mg of the title compound as the bis-trifluoroacetate salt mass spectrum (electrospray, m/e): M+H 542.0.

- 106 -

Example 1054-([1,3]Dioxolan-2-ylmethyl-methyl-amino)-but-2-enoic acid3-cyano-7-methoxy-quinolin-6-yl]-amide

In the manner of Example 103, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and [1,3]dioxolan-2-yl-methyl-methyl-amine was converted to 116.2 mg of the title compound as the bis-trifluoroacetate salt); mass spectrum (electrospray, m/e): M+H 526.0.

Example 1064-[Bis-(2-hydroxy-ethyl)-amino]-but-2-enoic acid[4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

In the manner of Example 103, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and bis-(2-hydroxy-ethyl) amine was converted to 22.2 mg of the title compound (free base), mass spectrum (electrospray, m/e): M+H 514.0 and 60.7 mg of the title compound as the bis-trifluoroacetate salt); mass spectrum (electrospray, m/e): M+H 514.0.

Example 1074-Thiomorpholin-4-yl-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

In the manner of Example 103, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and thiomorpholine was converted to 48.1 mg of the title compound (free base), mass spectrum (electrospray, m/e): M+H 512.0 and 33.2 mg of the title compound as the bis-trifluoroacetate salt); mass spectrum (electrospray, m/e): M+H 512.0.

Example 1084-[4-(2-Hydroxy-ethyl)-piperazin-1-yl]-but-2-enoic acid[4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

In the manner of Example 103, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and 4-(2-hydroxy-ethyl)

- 107 -

piperazine was converted to 32.3 mg of the title compound (free base) and mass spectrum (electrospray, m/e): M+H 539.1, 42.2 mg of the title compound as the bis-trifluoroacetate salt); mass spectrum (electrospray, m/e): M+H 539.1.

Example 109

4-(1,4,7-Trioxa-10-aza-cyclododec-10-yl)-but-2-enoic acid

[4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

In the manner of Example 103, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide and 1,4,7-trioxa-10-aza cyclododecane was converted to 37.5 mg of the title compound (free base) and mass spectrum (electrospray, m/e): M+H 584.1, 17.1 mg of the title compound as the bis-trifluoroacetate salt); mass spectrum (electrospray, m/e): M+H 584.1.

Example 110

4-(Methoxy-methyl-amino)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

A mixture of 1g of the 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide (2.04 mmol, 1 equiv.), 1.2g of N,O-Dimethylhydroxylamine hydrochloride (12.25 mmol, 9 equiv.), and 1.5g of sodium bicarbonate (18.38 mmol, 9 equiv.) in DMF (10ml) was stirred at room temperature for 24 hours. Ethyl acetate was added to the reaction mixture, and the crude product was filtered. After flash chromatography (ethyl acetate:methanol:triethylamine 40:4:1), 0.486 g of the title compound was isolated (50.7% yield); mp 210-217 °C.

Example 111

4-(4-Hydroxy-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

A mixture of 250 mg (0.51 mmol) of the 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide, and 103.2 mg (1.02 mmol) of 4-hydroxypiperidine in 2.25 ml dimethylformamide was stirred at room

- 108 -

temperature for 3 hr. Saturated sodium bicarbonate was added and the precipitate was filtered and washed with hexane to give the first crop of product. The filtrate was extracted with ethyl acetate and the organic layer with purified by preparative TLC to yield the second crop. The two crops were combined to give 105.8 mg (41%) tan solid: mp >215°C.

Example 112

4-[1,4'Bipiperidinyl-1'-yl-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

A mixture of 250 mg (0.51 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide, and 172 mg (1.02 mmol) of 4-piperidinopiperidine in 5.0 ml dimethylformamide was stirred at room temperature for 4 hr and at 60°C for 1 hr. After the mixture was cooled, the suspension was diluted with saturated sodium bicarbonate solution and extracted with ethyl acetate. The extracts were evaporated to an oil and purified by preparative TLC to yield 100 mg (40%) yellow solid: mp 140-144°C.

Example 113

4-Thiazolidin-3-yl-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

A mixture of 250 mg (0.51 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide, and 80µL (1.02 mmol) of thiazolidine in 2.25 ml dimethylformamide was stirred at room temperature for 19.5 hr. After the mixture was cooled, the suspension was diluted with saturated sodium bicarbonate solution and extracted with ethyl acetate. The extracts were evaporated to an oil and purified by preparative TLC to yield 95.6 mg (38%) yellow solid: mp 135-138°C.

- 109 -

Example 114**4-(2,6-Dimethyl-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide**

A mixture of 250 mg (0.51 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide, and 137 μ L (1.02 mmol) of cis-2,6-dimethylpiperidine in 2.25 ml dimethylformamide was stirred at room temperature for 3 hr. The reaction mixture was diluted with saturated sodium bicarbonate solution and extracted with ethyl acetate. The extracts were evaporated to an oil, washed with hexane and dried under reduced pressure to yield 170.4 mg (64%) tan solid: mp 120-122°C.

Example 115**4-[Bis-(2-hydroxy-propyl)-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide**

A mixture of 250 mg (0.51 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide, and 136 mg (1.02 mmol) of 1,1'-iminodi-2-propanol in 2.25 ml dimethylformamide was stirred at room temperature for 3 hr and at 60°C for 2 hr. After the mixture was cooled, saturated sodium bicarbonate solution was added and the solution was subsequently extracted with ethyl acetate. The extracts were evaporated to an oil, washed with hexane and dried under reduced pressure. Yield 240.1 mg (87%) tan solid: mp 122-125°C.

Example 116**4-(3-Hydroxy-pyrrolidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide**

A mixture of 250 mg (0.51 mmol) of 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide, and 85 μ L (1.02 mmol) of R-(+)-3-pyrrolidinol in 5.0 ml dimethylformamide was stirred at room temperature for 4 hr and at 60°C for 1 hr. After the mixture was cooled, saturated sodium bicarbonate solution was added and the solution was subsequently extracted

- 110 -

with ethyl acetate. The extracts were evaporated to an oil, and purified by preparative TLC. Yield 84.2 mg (33%) yellow solid: mp 215-220°C.

Example 117

4-[(2-Hydroxy-ethyl)-methyl-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide (250mg, .51mmol.) and 2-(methylamino)-ethanol(97 mg, 1.02mmol.) were dissolved with stirring, under nitrogen, in 9 ml anhydrous dimethylformamide. After 48 hours, the mixture was partitioned between saturated sodium bicarbonate solution, and ethyl acetate. The organic phase was separated, dried over magnesium sulfate, filtered, and evaporated to give a gum. It was chromatographed on silica gel, and eluted with 40/4/1 (ethyl acetate/methanol/triethylamine) producing 183 mg (74%) of the purified product as a yellow solid: mp 210-214°C.

Example 118

4-(2,5-Dimethyl-pyrrolidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

In the same manner as Example 117, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide (.51mmol.) was reacted with 2,5-dimethylpyrrolidine (1.02mmol., 101mg.) in dimethylformamide. The crude product was also purified via chromatography as Example 117, leaving 214mg. (82%) of the yellow product: mp 110-113°C.

Example 119

4-(4,4-Dihydroxy-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide

After the procedure of Example 117, 4-bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide (.51mmol.) was stirred in dimethylformamide with 4-piperidone monohydrate hydrochloride (470 mg.,

- 111 -

3.06mmol.), and sodium bicarbonate (386mg., 4.59mmol.), for 24 hours. The crude product was purified in the same manner as Example 117 and produced 192 mg. (72%) of the product as a yellow solid: mp 225-30°C.

Example 120

6-(4-Chlorobutylamino)-4-(3-chloro-4-fluorophenyamino)-7-methoxy -3-quinolinecarbonitrile

To a solution of 1.12 g of 4-chlorobutanal and 5.3 ml of 3M sulfuric acid in 11 ml of tetrahydrofuran, at 0°C, was added an solution of 2.0 g of 6-amino-4-(3-chloro-4-fluoro-phenylamino)-7-methoxy -3-quinolinecarbonitrile in 40 ml of dimethylformamide. To this was added portionwise 0.4 g of sodium borohydride. After 1 hr another 1.0 g of aldehyde, 10 ml of dimethylformamide, and 5 ml of 3M sulfuric acid was added followed by the portionwise addition of 0.8 g of sodium borohydride. After 2 hrs, the mixture was poured into water and the Ph was adjusted to 9. The mixture was extracted with ethyl acetate several times. The organic solution was dried over magnesium sulfate and the solvent was removed giving the title compound as an oil which was used without additional purification.

Example 121

4-(3-Chloro-4-fluoro-phenylamino)-7-methoxy-6-(4-morpholin-4-yl-butylamino)-quinoline-3-carbonitrile

Example 122

4-(3-Chloro-4-fluoro-phenylamino)-7-methoxy-6-pyrrolidin-1-yl-quinoline-3-carbonitrile

A mixture of 2.5 g of 6-(4-chlorobutylamino)-4-(3-chloro-4-fluorophenyamino)-7-methoxy -3-quinolinecarbonitrile, 7.54 g of morpholine, and 0.17 g of sodium iodide in 30 ml of dimethylformamide was stirred at 750°C for 7 hrs. The mixture was poured into dilute sodium bicarbonate and solid was collected. This material was dissolved in ethylacetate. The solution was dried over magnesium sulfate. The solvent was removed and the residue was chromatographed on silica gel using ethyl acetate-

5

- 112 -

10

15

methanol-triethylamine mixtures. A more polar component (0.54 g) was 4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-6-(4-morpholin-4-yl-butylamino)-quinoline-3-carbonitrile; a less polar component (0.28 g) is the compound of this invention, 4-(3-chloro-4-fluoro-phenylamino)-7-methoxy-6-pyrrolidin-1-yl-quinoline-3-carbonitrile, obtained as a yellow solid: mass spectrum electrospray, m/e): M+H 397.4.

Example 123

20

25

30

4-(3-Chloro-4-fluoroanilino)-7-methoxy-6-(1H-pyrrol-1-yl)-3-quinolinecarbonitrile
A dimethylformamide (4.5 ml) suspension of 6-amino-4-[(3-chloro-4-fluoroanilino)-7-methoxy-3-quinolinecarbonitrile (0.2 g, 0.5839 mmol), 2,5-dimethoxytetrahydrofuran (0.1 ml, 0.77 mmol) and 4-chloropyridinium chloride (0.05g, 0.333 mmol) was heated at 108 °C overnight. The reaction solution was mixed with saturated sodium bicarbonated solution, brine and ethyl acetate. The ethyl acetate layer was separated, filtered through silica gel, and dried up to give 119 mg of 4-(3-Chloro-4-fluoroanilino)-7-methoxy-6-(1H-pyrrol-1-yl)-3-quinolinecarbonitrile as a cream-colored solid, mp 192.5-193.5 °C: high resolution mass spectrometry (electrospray, m/e): M+H 393.0913.

35

20

Example 124

40

45

50

55

6-[(2-chloroethyl)amino]-4-(3-chloro-4-fluoroanilino)-7-methoxy-3-quinolinecarbonitrile

To a solution of 1 g of 6-amino-4-[(3-chloro-4-fluoroanilino)-7-methoxy-3-quinolinecarbonitrile (2.92 mmol) in 20 ml of dimethylformamide at 0 °C, was added a solution of 50% aqueous chloroformaldehyde (0.75 ml, 5.84 mmol) and 3 M sulfuric acid (2.92 ml, 8.76 mmol) in 5.3 ml of tetrahydrofuran, followed by portionwise addition of 1.1 g of sodium borohydride powder (29.68 mmol). After the reaction mixture was stirred overnight at room temperature, the precipitated product was collected and partitioned between ethyl acetate and 10 N sodium hydroxide solution. The organic layer was washed with brine and dried over sodium sulfate. The solvent was removed to yield 0.7648 g of a yellow solid. After recrystallization

- 113 -

from acetonitrile, 0.5831 g of bright needle crystals were obtained; mp 207.2-207.8 °C; high resolution mass spectrometry (EI, m/e): 404.060.

The filtrate was treated with 10 N sodium hydroxide and ethyl acetate, and worked up as above to afford additional 0.38 g of crude product.

Example 125

6-(1-Aziridinyl)-4-(3-chloro-4-fluoroanilino)-7-methoxy-3-quinolinecarbonitrile

To a solution of 0.202 g (0.5 mmol) of 6-[(2-chloroethyl)amino]-4-(3-chloro-4-fluoroanilino)-7-methoxy-3-quinolinecarbonitrile in 4 ml of dimethylformamide, were added 0.075 g of sodium iodide (0.5 mmol) and 0.069 g of potassium carbonate (0.5 mmol). After the reaction solution was heated at 75 °C overnight, it was decanted into cold saturated sodium bicarbonate solution. The organic layer was separated and dried over sodium sulfate. The solvent was removed to yield 0.19 g of a light brown solid. The crude product was washed with hexane to give 0.1g of a yellow solid, mp 197-199 °C; high resolution mass spectrometry (EI, m/e): M 368.0888.

Example 126

1-(Dimethylaminomethyleneamino)-3-chlorobenzene

A mixture of 3-chloroaniline (63.8 g, 0.50 mol) and dimethylformamide dimethyl acetal (106 ml, 0.75 mol) was heated at 100° for 2 h and evaporated at 60° at 0.5 mm Hg to give 91.8 g amber oil; ms 183.0 (M+H)•.

Example 127

1-(Dimethylaminomethyleneamino)-3-chloro-4-nitrobenzene

To a stirred solution of 1-(dimethylaminomethyleneamino)-3-chlorobenzene (67.5 g, 0.37 mol) in 148 ml of gl HOAc was added 70% nitric acid (70 ml, 1.11 mol) during 15 m with cooling at 10°. To the resulting solution was added Ac₂O (222 ml, 2.58 mol) during 30 m with cooling to maintain 15-20°. The solution was heated to 65° during 20 m. The resulting exothermic reaction was moderated with cool water at 65-68° for 45 m and then the reaction was heated at 65° for 90 m. The reaction mixture was cooled to 10°, stirred with DCM, and quenched with ice and 10 N NaOH (850

5

- 114 -

10

ml). The organic layer was separated, washed thoroughly with water, dried, filtered through Magnesol, and concentrated to give 62.2 g red gummy solid. Flash chromatography of the residue on silica gel with 20:4:1 DCM-EtOAc-MeOH gave an amber solid, mp 78-90°; ms 228.1 (M+H)⁺.

15

5

Example 128

(E/Z)-2-Cyano-3-(3-chloro-4-nitrophenylamino)acrylic Acid, Ethyl Ester

20

To a stirred mixture of 1-(dimethylaminomethyleneamino)-3-chloro-4-nitrobenzene (7.9 g, 35 mmol) and 17.4 ml of HOAc at 25° was added ethyl cyanoacetate (5.2 g, 46 mmol). The resulting mixture was refluxed for 1.5 h, cooled, and stirred in water for 45 m. The resulting amber solid was filtered off, washed with water followed by 5:1 hexane-EtOAc, and dried; mp 195-205°; ms 294.1(M-H).

25

Example 129

15

1,4-Dihydroquinoline-7-chloro-6-nitro-4-oxo-3-carbonitrile

30

A stirred mixture of (E/Z)-2-cyano-3-(3-chloro-4-nitrophenylamino)acrylic acid, ethyl ester (2.36 g, 8.0 mmol) and 240 ml of Dowtherm A was heated at 260°C for 2 h, cooled, diluted with hexane, and filtered. The tan solid thus obtained was digested with boiling EtOAc, filtered, and dried to give 1.47 g, mp 320-330° (dec); ms 248.1 (M-H).

35

20

Example 130

40

4,7-Dichloro-6-nitro-3-quinolinecarbonitrile

45

A stirred mixture of 1,4-dihydroquinoline-7-chloro-6-nitro-4-oxo-3-carbonitrile (14.7 g, 58.9 mmol) and 59 ml of phosphorous oxychloride was refluxed for 3 h. The phosphorous oxychloride was removed in vacuo, and the residue was stirred with methylene chloride at 0°C and treated with a slurry of ice and potassium carbonate. The mixture was filtered through Celite, and the organic layer of the filtrate was separated, washed with water, dried, and concentrated to give 10.7 g of tan solid. Recrystallization from hexane-DCM gave mp 143-153°; ms 266.7 (M-H).

50

30

55

5

- 115 -

Example 131

10

4-(3-Chloro-4-fluoroanilino)-7-chloro-6-nitro-3-quinolinecarbonitrile

15

A stirred mixture of 4,7-dichloro-6-nitro-3-quinolinecarbonitrile (10.7 g, 40 mmol), 3-chloro-4-fluoroaniline (7.0 g, 48 mmol), pyridine hydrochloride (4.6 g, 40 mmol), and 200 ml of 2-propanol was heated to reflux temperature and maintained for 1 h. The 2-propanol was evaporated off, and the residue was stirred in water with potassium bicarbonate (pH=8). The resulting solid was filtered, washed with water and 5:1 hexane-DCM, and dried. Recrystallization from EtOH gave 11.3 g of yellow solid, mp 259-263°; ms 377.1 (M+H)⁺.

20

10

Example 132

25

4-(3-Chloro-4-fluoroanilino)-7-(4-methyl-1-piperazinyl)-6-nitro-3-quinolinecarbonitrile

30

A stirred mixture of 4-(3-chloro-4-fluoroanilino)-7-chloro-6-nitro-3-quinolinecarbonitrile (1.88 g, 5.0 mmol), N-methylpiperazine (5 ml, 45 mmol), and 10 ml of toluene was refluxed for 45 m, evaporated to remove volatile matter, and stirred in water with potassium carbonate (2.75 g). The resulting solid was filtered, washed with water, and dried to give 2.26 g. An acetone solution was passed onto a pad of silica gel; elution with 50:2:1 acetone-MeOH-TEA and evaporation gave a red solid, mp 240-246°; ms 441.2 (M+H)⁺, 221.2 (M+2H)⁺2.

35

20

Example 133

40

6-Amino-4-(3-chloro-4-fluoroanilino)-7-(4-methyl-1-piperazinyl)-3-quinolinecarbonitrile

45

In the manner of Example 23 4-(3-chloro-4-fluoroanilino)-7-(4-methyl-1-piperazinyl)-6-nitro-3-quinolinecarbonitrile was reduced with iron powder and acetic acid in MeOH to give the title compound as an amorphous solid; ms 411.2 (M+H)⁺, 206.2 (M+2H)⁺2.

50

55

5

- 116 -

Example 134

10

N-[4-(3-Chloro-4-fluoroanilino)-3-cyano-7-(4-methyl-1-piperazinyl)-6-quinolinyl]-2-butynamide

15

20

25

To a stirred solution of 2-butyric acid (0.25 g, 3.0 mmol) in 1.5 ml of DCM at 0° was added DCC (0.21 g, 1.0 mmol). After 15 m the mixture was warmed to 25°, recooled to 0°, and treated with 6-amino-4-(3-chloro-4-fluoroanilino)-7-(4-methyl-1-piperazinyl)-3-quinolinecarbonitrile (0.21 g, 0.50 mmol) followed by 0.5 ml DCM rinse. The resulting mixture was stirred at 25° for 18 h and filtered to remove dicyclohexyl urea. The filtrate was partitioned with water which contained potassium carbonate (0.4 g, 3 mmol). The organic layer was washed with water, dried, and concentrated. The residue was passed as a solution in DCM onto a pad of silica gel. The product was eluted with 50:2:1 acetone-MeOH-TEA and concentrated to give 0.165 g amorphous solid; ms 477.2 (M+H)⁺, 239.1 (M+2H)²⁺.

15

Example 135

30

3-Chloro-N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]propanamide

35

40

45

To a stirred solution of 3-chloropropionic acid (0.65 g, 6.0 mmol) in 3 ml of DCM at 0° was added DCC (0.41 g, 2.0 mmol). After 15 m the mixture was warmed to 25°, recooled to 0°, and treated with 6-amino-4-(3-chloro-4-fluoroanilino)-7-(4-morpholinyl)-3-quinolinecarbonitrile (0.40 g, 1.0 mmol) followed by 1 ml DCM rinse. The resulting mixture was stirred at 25° for 20 h, diluted with DCM, and stirred with aqueous sodium bicarbonate. The mixture was filtered to remove dicyclohexyl urea. The organic layer of the filtrate was washed with water, dried, and concentrated. The residue was passed as a solution in DCM onto a pad of silica gel. The product was eluted with 25:25:2:1 DCM-EtOAc-MeOH-TEA and concentrated to give 0.38 g amorphous solid; ms 488.1 (M+H)⁺.

50

55

Example 136

N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]acrylamide

To a stirred solution of 3-chloro-N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]propanamide (0.30 g, 0.61 mmol) in 1.2 ml of THF at 0° was added 1.2 ml of 1.0 M KOtBu/tBuOH dropwise during 1 m. After 2 h at 0° the reaction was quenched with solid CO₂ and partitioned with DCM-water. The organic layer was washed with water, dried, and evaporated to give 0.28 g of the title compound as a white amorphous solid; ms 452.2 (M+H)⁺.

By using the methods described in Examples 1-136 above and the methods described in the patent applications WO-9843960 and WO-9909016, the compounds of this invention listed in Table 6 were prepared.

Table 6

Example	Compound	m.p. (°C)	mass spectrum
137	4-[(2-Methoxy-ethyl)-methyl-amino]-but-2-enoic acid [4-(3-bromo-phenylamino)-3-cyano-7-ethoxy-quinolin-6-yl]-amide	amorphous	538.0(M+H)
138	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(4-hydroxy-piperidin-1-yl)-propoxy]-6-methoxy-quinoline-3-carbonitrile	122-125	531.0(M+H)
139	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propoxy]-6-methoxy-quinoline-3-carbonitrile	133-137	560.1 (M+H)
140	4-(2-Bromo-4-chloro-phenylamino)-7-[2-[(2-hydroxy-ethyl)-methyl-amino]-ethoxy]-6-methoxy-quinoline-3-carbonitrile	186-188	597.0 (M+H), 254.2 (M+2H) ⁺
141	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-[(2-hydroxy-ethyl)-methyl-amino]-propoxy]-6-methoxy-quinoline-3-carbonitrile	129-131	533.0 (M+H)
142	4-(2,4-Dichloro-5-methoxy-phenylamino)-6-methoxy-7-(3-thiomorpholin-4-yl-propoxy)-quinoline-3-carbonitrile	116-118	505.2(M+H)
143	4-(2,4-Dichloro-5-methoxy-phenylamino)-6-methoxy-7-[3-(2-methoxy-ethylamino)-propoxy]-quinoline-3-carbonitrile	98-102	529.2 (M+H)
144	4-(2,4-Dichloro-5-methoxy-phenylamino)-6-methoxy-7-[3-(4-methyl-piperidin-1-yl)-propoxy]-quinoline-3-carbonitrile	114-117	587.2 (M+H)
145	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(2,6-dimethyl-morpholin-4-yl)-propoxy]-6-methoxy-quinoline-3-carbonitrile	155-157	545.3 (M+H)
146	4-(2-Bromo-4-chloro-phenylamino)-7-[2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-ethoxy]-6-methoxy-quinoline-3-carbonitrile	156-158	562.1 (M+H), 281.7 (M+2H) ⁺
147	4-(2-Bromo-4-chloro-phenylamino)-7-[2-(4-hydroxy-piperidin-1-yl)-ethoxy]-6-methoxy-quinoline-3-carbonitrile	165-167	533.1 (M+H), 267.1 (M+2H) ⁺
148	4-(2-Bromo-4-chloro-phenylamino)-6-methoxy-7-(2-thiomorpholin-4-yl-ethoxy)-quinoline-3-carbonitrile	164-166	533.0 (M+H), 268.1 (M+2H) ⁺

- 118 -

149	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(2,5-dimethyl-pyrrolidin-1-yl)-propoxy]-6-methoxy-quinoline-3-carbonitrile	115-120	529.2 (M+H)
150	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(3-hydroxy-propylamino)-propoxy]-6-methoxy-quinoline-3-carbonitrile	142-147	505.2 (M+H)
151	1-[3-[3-Cyano-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinolin-7-yloxy]-propyl]-piperidine-4-carboxylic acid ethyl ester	95-101	587.2 (M+H)
152	7-[3-(4-acetyl-1-piperazinyl)propoxy]-4-[(2,4-dichloro-5-methoxyphenyl)amino]-6-methoxy-3-quinolinecarbonitrile	115-118	558.2 (M+H)
153	4-(3-chloro-4-fluoroanilino)-7-methoxy-6-(4-morpholinyl)-3-quinolinecarbonitrile		413.2 (M+H)
154	7-[3-(4-Benzyl-piperazin-1-yl)-propoxy]-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinoline-3-carbonitrile	140-142	606.2 (M+H)
155	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(2-hydroxy-ethylamino)-propoxy]-6-methoxy-quinoline-3-carbonitrile	161-164	491.1 (M+H)
156	4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-[ethyl-(2-hydroxy-ethyl)-amino]-propoxy]-6-methoxy-quinoline-3-carbonitrile	162-165	519.2 (M+H)
157	7-[3-[Bis-(2-methoxy-ethyl)-amino]-propoxy]-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinoline-3-carbonitrile	112-113	563.1 (M+H)
158	7-[3-[Bis-(2-hydroxy-ethyl)-amino]-propoxy]-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinoline-3-carbonitrile	156-159	535.1 (M+H)
159	4-(3-chloro-4-fluoroanilino)-7-(4-morpholinyl)-6-nitro-3-quinolinecarbonitrile	235-239	428.1 (M+H)
160	N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]-2-butanamide	260-266d	464.1 (M+H)
161	6-amino-4-(3-chloro-4-fluoroanilino)-7-(4-morpholinyl)-3-quinolinecarbonitrile	amorphous	398.2 (M+H)
162	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-(3-[2-(4-morpholinyl)ethyl]amino)propoxy]-3-quinolinecarbonitrile	75-80	560.2 (M+H)
163	7-[3-[(2-anilinoethyl)amino]propoxy]-4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-3-quinolinecarbonitrile	90-94	566.2 (M+H)
164	N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]acrylamide	amorphous	
165	4-(3-chloro-4-fluoroanilino)-7-[4-[2-(dimethylamino)ethyl]-1-piperazinyl]-6-nitro-3-quinolinecarbonitrile		467.2 (M+H)
166	6-amino-4-(3-chloro-4-fluoroanilino)-7-[4-[2-(dimethylamino)ethyl]-1-piperazinyl]-3-quinolinecarbonitrile		468.2 (M+H), 234.7 (M+2H) ⁺¹
167	N-(4-(3-chloro-4-fluoroanilino)-3-cyano-7-[4-[2-(dimethylamino)ethyl]-1-piperazinyl]-6-quinolinyl)acrylamide		522.2 (M+H), 261.7 (M+2H) ⁺¹
168	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[(2-[4-(2-methoxyethyl)-1-piperazinyl]ethyl)amino]-3-quinolinecarbonitrile	53-55	559.3 (M+H), 280.2 (M+2H) ⁺²
169	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(2H-1,2,3-triazol-2-yl)propoxy]-3-quinolinecarbonitrile	190-191	499.4 (M+H)
170	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-1,2,3-triazol-1-yl)propoxy]-3-quinolinecarbonitrile	188-190	499.4 (M+H)

5

- 119 -

10

15

20

25

30

35

40

45

50

55

171	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-(3-thienyl)-3-quinolinecarbonitrile	215-218	456.3 (M+H)
172	4-[(E)-2-(2-quinolinyl)ethenyl]aniline	53-54	572.5 (M+H), 286.9 (M+2H) ¹²
173	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-([2-(2H-1,2,3-triazol-2-yl)ethyl]amino)-3-quinolinecarbonitrile	210-211	484.1 (M+H)
174	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-([2-(1H-1,2,3-triazol-1-yl)ethyl]amino)-3-quinolinecarbonitrile	225-228	484.1 (M+H)
175	4-(2,4-dichloro-5-methoxyanilino)-7-(3-thienyl)-3-quinolinecarbonitrile	211-212	426.0 (M+H)
176	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-1,2,4-triazol-1-yl)propoxy]-3-quinolinecarbonitrile	206-208	499.1 (M+H)
177	4-(2,4-dichloro-5-methoxyanilino)-7-[3-(1H-imidazol-1-yl)propoxy]-6-methoxy-3-quinolinecarbonitrile	155-170	498.1 (M+H), 249.6 (M+2H) ¹²
178	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-pyrazol-1-yl)propoxy]-3-quinolinecarbonitrile	187-188	498.1 (M+H)
179	N-[3-cyano-4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-quinolinyl]-N-[4-(4-ethyl-1-piperazinyl)butyl]acetamide	57	599.2 (M+H), 300.3 (M+2H) ¹²
180	N-[3-cyano-4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-quinolinyl]-N-(3-(4-ethyl-1-piperazinyl)propyl)acetamide	58.5-59	585.1 (M+H), 293.2 (M+2H) ¹²
181	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-[4-(2-methoxyethyl)-1-piperazinyl]propoxy]-3-quinolinecarbonitrile	118-120	574.1 (M+H)
182	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-(1H-pyrrol-1-yl)-3-quinolinecarbonitrile	229-230	439.1 (M+H)
183	4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(1H-1,2,3-triazol-1-yl)ethoxy]-3-quinolinecarbonitrile	180-182	483.0 (M+H)
184	4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(2H-1,2,3-triazol-2-yl)ethoxy]-3-quinolinecarbonitrile	93-103	483.0 (M+H)
185	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-tetrazol-1-yl)propoxy]-3-quinolinecarbonitrile	210-214	500.1 (M+H)
186	4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(2H-tetrazol-2-yl)propoxy]-3-quinolinecarbonitrile	228-230	500.0 (M+H)
187	4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(1H-1,2,3-triazol-1-yl)ethoxy]-3-quinolinecarbonitrile	180-184	483.0 (M+H)
188	4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(2H-1,2,3-triazol-2-yl)ethoxy]-3-quinolinecarbonitrile	95-103	483.0 (M+H)
189	4-(2,4-dichloro-5-methoxyanilino)-7-[3-([2-(dimethylamino)ethyl](methyl)amino)propoxy]-6-methoxy-3-quinolinecarbonitrile	85-90	532.1 (M+H), 266.7 (M+2H) ¹²

Claims

5

10

15

20

25

30

35

40

45


50

55

- 120 -

WHAT IS CLAIMED IS:

1. A compound of Formula 1 having the structure:



wherein:



X is cycloalkyl of 3 to 7 carbon atoms, which may be optionally substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a pyridinyl, pyrimidinyl, or phenyl ring wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, carboxyalkyl of 2-7 carbon atoms, carboalkoxyalkyl of 3-8 carbon atoms, aminoalkyl of 1-5 carbon atoms, N-alkylaminoalkyl of 2-9 carbon atoms, N,N-dialkylaminoalkyl of 3-10 carbon atoms, N-alkylaminoalkoxy of 2-9 carbon atoms, N,N-dialkylaminoalkoxy of 3-10 carbon atoms, mercapto, and benzoylamino;

Z is -NH-, -O-, -S-, or -NR-;

- 121 -



R is alkyl of 1-6 carbon atoms, or carboalkyl of 2-7 carbon atoms;

G₁, G₂, R₁, and R₄ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, alkenyloxy of 2-6 carbon atoms, alkynyloxy of 2-6 carbon atoms, hydroxymethyl, halomethyl, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkenoyloxymethyl of 4-9 carbon atoms, alkynoyloxymethyl of 4-9 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulphonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, alkenylsulfonamido of 2-6 carbon atoms, alkynylsulfonamido of 2-6 carbon atoms, hydroxy, trifluoromethyl, trifluoromethoxy, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, amino, hydroxyamino, alkoxyamino of 1-4 carbon atoms, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, phenylamino, benzylamino,




with the proviso that either G₁ or G₂ or both G₁ and G₂ must be a radical selected from the group

- 122 -



Y is a divalent radical selected from the group consisting of




R₇ is -NR₆R₆, -J, -OR₆, -N(R₆)₃⁺, or -NR₆(OR₆);

R'₇ is -NR₆(OR₆), -N(R₆)₃⁺, alkenoxy of 1-6 carbon atoms, alkynoxy of 1-6 carbon atoms, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, N-alkyl-N-alkynylamino of 4 to 12 carbon atoms, N-alkenyl-N-alkynylamino of 4 to 12 carbon atoms, or N,N-dialkynylamino of 6-12 carbon atoms with the proviso that the alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

M is >NR₆, -O-, >N-(C(R₆)₂)_pNR₆R₆, or >N-(C(R₆)₂)_p-OR₆;

W is >NR₆, -O- or is a bond;

Het is a heterocycle selected from the group consisting of morpholine, thiomorpholine, thiomorpholine S-oxide, thiomorpholine S,S-dioxide, piperidine, pyrrolidine, aziridine, pyridine, imidazole, 1,2,3-triazole, 1,2,4-triazole, thiazole, thiazolidine, tetrazole, piperazine, furan, thiophene, tetrahydrothiophene, tetrahydrofuran, dioxane, 1,3-dioxolane,



5

10

15

20

5

25

30

35

40

45

50

55

5

- 123 -

10

wherein the heterocycle is optionally mono- or di-substituted on carbon or nitrogen with R_6 , optionally mono- or di-substituted on carbon with hydroxy, $-N(R_6)_2$, or $-OR_6$, optionally mono or di-substituted on carbon with the mono-valent radicals $-(C(R_6)_2)_sOR_6$ or $-(C(R_6)_2)_sN(R_6)_2$, or optionally

15

5

mono or di-substituted on a saturated carbon with divalent radicals $-O-$ or $-O(C(R_6)_2)_sO-$;

20

R_6 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-

10

6 carbon atoms, cycloalkyl of 1-6 carbon atoms, carboalkyl of 2-7 carbon atoms, carboxyalkyl (2-7 carbon atoms), phenyl, or phenyl optionally substituted with one or more halogen, alkoxy of 1-6 carbon atoms, trifluoromethyl, amino, alkylamino of 1-3 carbon atoms, dialkylamino of 2-6 carbon atoms, nitro, cyano, azido, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, carboxyl, carboalkoxy of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, or alkyl of 1-6 carbon atoms;

25

30

15

35

R_2 , is selected from the group consisting of

20

40

45


50

55


5

- 124 -


10




15




20




25



30



35




40

45


50

55


- 125 -



R₃ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,






with the proviso that at least one of the R₃ groups is selected from the group



- 126 -

R₅ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,



5

R₈ and R₉ are each, independently, -(C(R₆)₂)_rNR₆R₆, or -(C(R₆)₂)_rOR₆;

J is independently hydrogen, chlorine, fluorine, or bromine;

Q is alkyl of 1-6 carbon atoms or hydrogen;

a = 0 or 1;

10 g = 1-6;

k = 0-4;

n is 0-1;

p = 2-4;

q = 0-4;

15 r = 1-4;

s = 1-6;

u = 0-4 and v = 0-4, wherein the sum of u+v is 2-4;

or a pharmaceutically acceptable salt thereof,

provided that

20 when R₆ is alkenyl of 2-7 carbon atoms or alkynyl of 2-7 carbon atoms, such alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

and further provided that

when Y is -NR₆- and R₇ is -NR₆R₆, -N(R₆)₃⁺, or -NR₆(OR₆), then g = 2-6;

25 when M is -O- and R₇ is -OR₆, then p = 1-4;

5

- 127 -

10

when Y is -NR₆-, then k = 2-4;

when Y is -O- and M or W is -O-, then k = 1-4;

when W is not a bond with Het bonded through a nitrogen atom, then q = 2-4;

and when W is a bond with Het bonded through a nitrogen atom and Y is -O-

15

5 or -NR₆-, then k = 2-4.

20

2. The compound according to claim 1 wherein Z is -NH- and n = 0 or a pharmaceutically acceptable salt thereof.

25

10 3. The compound according to claim 2 wherein X is optionally substituted phenyl or a pharmaceutically acceptable salt thereof.

15

4. The compound according to claim 3 wherein R₁ and R₄ are hydrogen or a pharmaceutically acceptable salt thereof.

30

5. The compound according to claim 1, which is:

35

a) 1-Methyl-1,2,5,6-tetrahydro-pyridine-3-carboxylic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

40

b) N-[4-[(3-Bromophenyl)amino]-3-cyano-6-quinolinyl]-4-(N-allyl-N-methylamino)-2-butyramide or a pharmaceutically acceptable salt thereof;

45

25 c) N-[4-[(3-Bromophenyl)amino]-3-cyano-6-quinolinyl]-4-(N-methoxyethyl-N-methylamino)-2-butyramide or a pharmaceutically acceptable salt thereof;

50

30

d) N-[4-[(3-Bromophenyl)amino]-3-cyano-6-quinolinyl]-4-(bis-(2-methoxy-ethyl)amino)-2-butyramide or a pharmaceutically acceptable salt thereof;

55

5

- 128 -

10

e) 4-Methoxymethoxy-but-2-ynoic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

15

5 f) 4-(4-Chloro-2-fluoro-phenylamino)-6-methoxy-7-(2-pyridin-4-yl-ethoxy)-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

20

g) 4-(2-Methoxy-ethoxy)-but-2-ynoic acid [4-(3-bromo-phenylamino)-3-cyano-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

10

h) 4-((2S)-2-Methoxymethylpyrrolidin-1-yl)but-2-ynoic Acid [4-(3-bromophenylamino)-3-cyanoquinolin-6-yl]amide or a pharmaceutically acceptable salt thereof;

25

15 i) 4-(1,4-Dioxo-8-azaspiro[4,5]dec-8-yl)but-2-ynoic Acid [4-(3-Bromophenylamino)-3-cyanoquinolin-6-yl] amide or a pharmaceutically acceptable salt thereof;

30

j) 4-(3-Bromo-phenylamino)-6-(2-ethoxy-3,4-dioxo-cyclobut-1-enylamino)-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

35

20

k) 4-[(2-Methoxy-ethyl)-methyl-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

40

25 l) (S)-4-(2-Methoxymethyl-pyrrolidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide dihydrochloride or a pharmaceutically acceptable salt thereof;

45

50 30 m) 4-(3-Hydroxymethyl-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

55

5

- 129 -

10

n) 4-(1,4-Dioxo-8-aza-spiro[4.5]dec-8-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

15

5

o) 4-(2-Hydroxymethyl-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

20

10

p) 4-Bromo-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

25

q) 4-(3-hydroxy-4-methyl-phenylamino)-6-methoxy-7-(3-pyridin-4-yl-propoxy)-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

15

30

r) 4-Diallylamino-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

35

20

s) 4-[Bis-(2-methoxy-ethyl)-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

40

t) 4-([1,3]Dioxolan-2-ylmethyl-methyl-amino)-but-2-enoic acid [4-(3-cyano-7-methoxy-quinolin-6-yl)-amide or a pharmaceutically acceptable salt thereof;

25

45

u) 4-[Bis-(2-hydroxy-ethyl)-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

50

30

55

5

- 130 -

10

v) 4-Thiomorpholin-4-yl-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

15

5 w) 4-[4-(2-Hydroxy-ethyl)-piperazin-1-yl]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

20

10 x) 4-(1,4,7-Trioxa-10-aza-cyclododec-10-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

25

15 y) 4-(Methoxy-methyl-amino)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

30

z) 4-(4-Hydroxy-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

35

20

aa) 4-[1,4']Bipiperidinyl-1'-yl-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

40

25 bb) 4-Thiazolidin-3-yl-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

45

50 30 cc) 3-{3-[4-(3-Chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]carbamoyl}-allyl}-4-methyl-thiazol-3-ium bromide or a pharmaceutically acceptable salt thereof;

55

5

- 131 -

10

dd) 4-(2,6-Dimethyl-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

15

5

ee) 4-[Bis-(2-hydroxy-propyl)-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

20

10

ff) 4-(3-Hydroxy-pyrrolidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

25

gg) 4-[(2-Hydroxy-ethyl)-methyl-amino]-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

30

15

hh) 4-(2,5-Dimethyl-pyrrolidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

35

20

ii) 4-(4,4-Dihydroxy-piperidin-1-yl)-but-2-enoic acid [4-(3-chloro-4-fluoro-phenylamino)-3-cyano-7-methoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

40

25

jj) 4-(3-Chloro-4-fluoro-phenylamino)-7-methoxy-6-pyrrolidin-1-yl-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

45

kk) 4-(3-Chloro-4-fluoroanilino)-7-methoxy-6-(1H-pyrrol-1-yl)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

50

30

55

5

- 132 -

10

ll) 6-(1-Aziridinyl)-4-(3-chloro-4-fluoroanilino)-7-methoxy-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

15

5 mm) 4-[(2-Methoxy-ethyl)-methyl-amino]-but-2-enoic acid [4-(3-bromophenylamino)-3-cyano-7-ethoxy-quinolin-6-yl]-amide or a pharmaceutically acceptable salt thereof;

20

nn) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(4-hydroxy-piperidin-1-yl)-propoxy]-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

10

25

oo) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-{3-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-propoxy}-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

15

30

pp) 4-(2-Bromo-4-chloro-phenylamino)-7-{2-[(2-hydroxy-ethyl)-methyl-amino]-ethoxy}-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

35

20 qq) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-{3-[(2-hydroxy-ethyl)-methyl-amino]-propoxy}-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

40

rr) 4-(2,4-Dichloro-5-methoxy-phenylamino)-6-methoxy-7-(3-thiomorpholin-4-yl-propoxy)-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

25

45

ss) 4-(2,4-Dichloro-5-methoxy-phenylamino)-6-methoxy-7-[3-(2-methoxy-ethylamino)-propoxy]-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

50

30

55

5

- 133 -

10

tt) 4-(2,4-Dichloro-5-methoxy-phenylamino)-6-methoxy-7-[3-(4-methyl-piperidin-1-yl)-propoxy]-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

15

5 uu) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(2,6-dimethyl-morpholin-4-yl)-propoxy]-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

20

10 vv) 4-(2-Bromo-4-chloro-phenylamino)-7-{2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-ethoxy}-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

25

15 ww) 4-(2-Bromo-4-chloro-phenylamino)-7-[2-(4-hydroxy-piperidin-1-yl)-ethoxy]-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

30

xx) 4-(2-Bromo-4-chloro-phenylamino)-6-methoxy-7-(2-thiomorpholin-4-yl)-ethoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

35

20 yy) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(2,5-dimethyl-pyrrolidin-1-yl)-propoxy]-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

40

25 zz) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(3-hydroxy-propylamino)-propoxy]-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

45

50 30 aaa) 1-{3-[3-Cyano-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinolin-7-yloxy]-propyl}-piperidine-4-carboxylic acid ethyl ester or a pharmaceutically acceptable salt thereof;

55

5

- 134 -

10

bbb) 7-[3-(4-acetyl-1-piperazinyl)propoxy]-4-[(2,4-dichloro-5-methoxyphenyl)amino]-6-methoxy-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

15

5 ccc) 4-(3-chloro-4-fluoroanilino)-7-methoxy-6-(4-morpholinyl)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

20

ddd) 7-[3-(4-Benzyl-piperazin-1-yl)-propoxy]-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

25

eee) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-[3-(2-hydroxy-ethylamino)-propoxy]-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

15

30

fff) 4-(2,4-Dichloro-5-methoxy-phenylamino)-7-{3-[ethyl-(2-hydroxy-ethyl)-amino]-propoxy}-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

35

20 ggg) 7-{3-[Bis-(2-methoxy-ethyl)-amino]-propoxy}-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

40

hhh) 7-{3-[Bis-(2-hydroxy-ethyl)-amino]-propoxy}-4-(2,4-dichloro-5-methoxy-phenylamino)-6-methoxy-quinoline-3-carbonitrile or a pharmaceutically acceptable salt thereof;

45

iii) 4-(3-chloro-4-fluoroanilino)-7-(4-morpholinyl)-6-nitro-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

50

30

55

5

- 135 -

10

jjj) N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]-2-butynamide or a pharmaceutically acceptable salt thereof;

15

5 kkk) 6-amino-4-(3-chloro-4-fluoroanilino)-7-(4-morpholinyl)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

20

lll) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-(3-{[2-(4--morpholinyl)ethyl]amino}propoxy)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

10

25

mmm) 7-{3-[(2-anilinoethyl)amino]propoxy}-4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

30

15 nnn) N-[4-(3-chloro-4-fluoroanilino)-3-cyano-7-(4-morpholinyl)-6-quinolinyl]acrylamide or a pharmaceutically acceptable salt thereof;

35

ooo) 4-(3-chloro-4-fluoroanilino)-7-{4-[2-(dimethylamino)ethyl]-1-piperazinyl}-6-nitro-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

40

20 ppp) 6-amino-4-(3-chloro-4-fluoroanilino)-7-{4-[2-(dimethylamino)ethyl]-1-piperazinyl}-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

45

25 qqq) N-(4-(3-chloro-4-fluoroanilino)-3-cyano-7-{4-[2-(dimethylamino)ethyl]-1-piperazinyl}-6-quinolinyl)acrylamide or a pharmaceutically acceptable salt thereof;

50

30 rrr) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-({2-[4-(2-methoxyethyl)-1-piperazinyl]ethyl}amino)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

55

5

- 136 -

10

sss) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(2H-1,2,3-triazol-2-yl)propoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

15

5 ttt) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-1,2,3-triazol-1-yl)propoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

20

uuu) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-(3-thienyl)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

25

vvv) 4-[(E)-2-(2-quinolinyl)ethenyl]aniline or a pharmaceutically acceptable salt thereof;

30

15 www) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[[2-(2H-1,2,3-triazol-2-yl)ethyl]amino]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

35

20 xxx) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[[2-(1H-1,2,3-triazol-1-yl)ethyl]amino]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

40

yyy) 4-(2,4-dichloro-5-methoxyanilino)-7-(3-thienyl)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

25

45

zzz) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-1,2,4-triazol-1-yl)propoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

50

30 aaaa) 4-(2,4-dichloro-5-methoxyanilino)-7-[3-(1H-imidazol-1-yl)propoxy]-6-methoxy-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

55

5

- 137 -

10

bbbb) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-pyrazol-1-yl)propoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

15

5

cccc) N-[3-cyano-4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-quinoliny]-N-[4-(4-ethyl-1-piperazinyl)butyl]acetamide or a pharmaceutically acceptable salt thereof;

20

10

dddd) N-[3-cyano-4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-quinoliny]-N-(3-(4-ethyl-1-piperazinyl)propyl)acetamide or a pharmaceutically acceptable salt thereof;

25

eeee) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-{3-[4-(2-methoxyethyl)-1-piperazinyl]propoxy}-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

30

15

ffff) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-(1H-pyrrol-1-yl)-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

35

20

gggg) 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(1H-1,2,3-triazol-1-yl)ethoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

40

hhhh) 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(2H-1,2,3-triazol-2-yl)ethoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

25

45

iiii) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(1H-tetrazol-1-yl)propoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

50

30

55

- 138 -


iiii) 4-(2,4-dichloro-5-methoxyanilino)-6-methoxy-7-[3-(2H-tetrazol-2-yl)propoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

kkkk) 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(1H-1,2,3-triazol-1-yl)ethoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof;

llll) 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-[2-(2H-1,2,3-triazol-2-yl)ethoxy]-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof; or

mmmm) 4-(2,4-dichloro-5-methoxyanilino)-7-{3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy}-6-methoxy-3-quinolinecarbonitrile or a pharmaceutically acceptable salt thereof.

6. A method of treating, inhibiting the growth of, or eradicating a neoplasm in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of formula 1 having the structure



wherein:

X is cycloalkyl of 3 to 7 carbon atoms, which may be optionally substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a pyridinyl, pyrimidinyl, or phenyl ring wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon

5

- 139 -

10

15

20

25

30

35

40

45

50

55

atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, carboxyalkyl of 2-7 carbon atoms, carboalkoxyalkyl of 3-8 carbon atoms, aminoalkyl of 1-5 carbon atoms, N-alkylaminoalkyl of 2-9 carbon atoms, N,N-dialkylaminoalkyl of 3-10 carbon atoms, N-alkylaminoalkoxy of 2-9 carbon atoms, N,N-dialkylaminoalkoxy of 3-10 carbon atoms, mercapto, and benzoylamino;



Z is -NH-, -O-, -S-, or -NR- ;

R is alkyl of 1-6 carbon atoms, or carboalkyl of 2-7 carbon atoms;




G₁, G₂, R₁, and R₄ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, alkenyloxy of 2-6 carbon atoms, alkynyloxy of 2-6 carbon atoms, hydroxymethyl, halomethyl, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkenoyloxymethyl of 4-9 carbon atoms, alkynoyloxymethyl of 4-9 carbon atoms, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulphonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, alkenylsulfonamido of 2-6 carbon atoms, alkynylsulfonamido of 2-6 carbon atoms, hydroxy, trifluoromethyl, trifluoromethoxy, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, amino, hydroxyamino, alkoxyamino of 1-4 carbon atoms, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, N-alkylcarbamoyle, N,N-dialkylcarbamoyle, N-alkyl-N-

- 140 -


alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, phenylamino, benzylamino,



with the proviso that either G₁ or G₂ or both G₁ and G₂ must be a radical selected from the group



Y is a divalent radical selected from the group consisting of



R₇ is -NR₆R₆, -J, -OR₆, -N(R₆)₃⁺, or -NR₆(OR₆);


R₇ is -NR₆(OR₆), -N(R₆)₃⁺, alkenoxy of 1-6 carbon atoms, alkynoxy of 1-6 carbon atoms, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, N-alkyl-N-alkynylamino of 4 to 12 carbon atoms, N-alkenyl-N-alkynylamino of 4 to 12 carbon atoms, or N,N-dialkynylamino of 6-12 carbon atoms with the proviso that the alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

M is >NR₆, -O-, >N-(C(R₆)₂)_pNR₆R₆, or >N-(C(R₆)₂)_p-OR₆;

- 141 -

W is $>NR_6$, -O- or is a bond;

Het is a heterocycle selected from the group consisting of morpholine, thiomorpholine, thiomorpholine S-oxide, thiomorpholine S,S-dioxide, piperidine, pyrrolidine, aziridine, pyridine, imidazole, 1,2,3-triazole, 1,2,4-triazole, thiazole, thiazolidine, tetrazole, piperazine, furan, thiophene, tetrahydrothiophene, tetrahydrofuran, dioxane, 1,3-dioxolane,










tetrahydropyran, and

wherein the heterocycle is optionally mono- or di-substituted on carbon or nitrogen with R_6 , optionally mono- or di-substituted on carbon with hydroxy, $-N(R_6)_2$, or $-OR_6$, optionally mono or di-substituted on carbon with the mono-valent radicals $-(C(R_6)_2)_sOR_6$ or $-(C(R_6)_2)_sN(R_6)_2$, or optionally mono or di-substituted on a saturated carbon with divalent radicals -O- or $-O(C(R_6)_2)_sO-$;

R_6 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 1-6 carbon atoms, carboalkyl of 2-7 carbon atoms, carboxyalkyl (2-7 carbon atoms), phenyl, or phenyl optionally substituted with one or more halogen, alkoxy of 1-6 carbon atoms, trifluoromethyl, amino, alkylamino of 1-3 carbon atoms, dialkylamino of 2-6 carbon atoms, nitro, cyano, azido, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, carboxyl, carboalkoxy of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, or alkyl of 1-6 carbon atoms;

R_2 , is selected from the group consisting of

- 142 -



5

10

15

20

25

30

35


40

45


50

55


- 143 -



R₃ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,






with the proviso that at least one of the R₃ groups is selected from the group



- 144 -

R₅ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,



R₈, and R₉ are each, independently, -(C(R₆)₂)_rNR₆R₆, or -(C(R₆)₂)_rOR₆;

J is independently hydrogen, chlorine, fluorine, or bromine;

Q is alkyl of 1-6 carbon atoms or hydrogen;

a = 0 or 1;

g = 1-6;

k = 0-4;

n is 0-1;

p = 2-4;

q = 0-4;

r = 1-4;

s = 1-6;

u = 0-4 and v = 0-4, wherein the sum of u+v is 2-4;

or a pharmaceutically acceptable salt thereof,

provided that

when R₆ is alkenyl of 2-7 carbon atoms or alkynyl of 2-7 carbon atoms, such alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

and further provided that

when Y is -NR₆- and R₇ is -NR₆R₆, -N(R₆)₃⁺, or -NR₆(OR₆), then g = 2-6;

when M is -O- and R₇ is -OR₆, then p = 1-4;

- 145 -

when Y is -NR₆-, then k = 2-4;


when Y is -O- and M or W is -O-, then k = 1-4;

when W is not a bond with Het bonded through a nitrogen atom, then q = 2-4;

and when W is a bond with Het bonded through a nitrogen atom and Y is -O- or -NR₆-, then k = 2-4.

7. The method according to claim 6 wherein the neoplasm is selected from the group consisting of breast, kidney, bladder, mouth, larynx, esophagus, stomach, colon, ovary, and lung.

8. A method of treating, inhibiting the progression of, or eradicating polycystic kidney disease in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of formula 1 having the structure



wherein:

X is cycloalkyl of 3 to 7 carbon atoms, which may be optionally substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a pyridinyl, pyrimidinyl, or phenyl ring wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon

5

- 146 -

10

15

20

atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, carboxyalkyl of 2-7 carbon atoms, carboalkoxyalkyl of 3-8 carbon atoms, aminoalkyl of 1-5 carbon atoms, N-alkylaminoalkyl of 2-9 carbon atoms, N,N-dialkylaminoalkyl of 3-10 carbon atoms, N-alkylaminoalkoxy of 2-9 carbon atoms, N,N-dialkylaminoalkoxy of 3-10 carbon atoms, mercapto, and benzoylamino;

10 Z is -NH-, -O-, -S-, or -NR- ;

R is alkyl of 1-6 carbon atoms, or carboalkyl of 2-7 carbon atoms;

25

G₁, G₂, R₁, and R₄ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon

30

15

atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, alkenyloxy of 2-6 carbon atoms, alkynyloxy of 2-6 carbon atoms, hydroxymethyl, halomethyl, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkenoyloxymethyl of 4-9 carbon atoms, alkynoyloxymethyl of 4-9 carbon atoms, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulphonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, alkenylsulfonamido of 2-6 carbon atoms, alkynylsulfonamido of 2-6 carbon atoms, hydroxy, trifluoromethyl, trifluoromethoxy, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, amino, hydroxyamino, alkoxyamino of 1-4 carbon atoms, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, phenylamino, benzylamino,

35

20

40



25

45



50

55


- 147 -



with the proviso that either G₁ or G₂ or both G₁ and G₂ must be a radical selected from the group



Y is a divalent radical selected from the group consisting of



10

R₇ is -NR₆R₆, -J, -OR₆, -N(R₆)₃⁺, or -NR₆(OR₆);

R'₇ is -NR₆(OR₆), -N(R₆)₃⁺, alkenoxy of 1-6 carbon atoms, alkynoxy of 1-6 carbon atoms, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, N-alkyl-N-alkynylamino of 4 to 12 carbon atoms, N-alkenyl-N-alkynylamino of 4 to 12 carbon atoms, or N,N-dialkynylamino of 6-12 carbon atoms with the proviso that the alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

M is >NR₆, -O-, >N-(C(R₆)₂)_pNR₆R₆, or >N-(C(R₆)₂)_p-OR₆;


W is >NR₆, -O- or is a bond;

20

Het is a heterocycle selected from the group consisting of morpholine, thiomorpholine, thiomorpholine S-oxide, thiomorpholine S,S-dioxide,

- 148 -

piperidine, pyrrolidine, aziridine, pyridine, imidazole, 1,2,3-triazole, 1,2,4-triazole, thiazole, thiazolidine, tetrazole, piperazine, furan, thiophene, tetrahydrothiophene, tetrahydrofuran, dioxane, 1,3-dioxolane ,



tetrahydropyran, and ;

wherein the heterocycle is optionally mono- or di-substituted on carbon or nitrogen with R_6 , optionally mono- or di-substituted on carbon with hydroxy, $-\text{N}(\text{R}_6)_2$, or $-\text{OR}_6$, optionally mono or di-substituted on carbon with the mono-valent radicals $-(\text{C}(\text{R}_6)_2)_s\text{OR}_6$ or $-(\text{C}(\text{R}_6)_2)_s\text{N}(\text{R}_6)_2$, or optionally mono or di-substituted on a saturated carbon with divalent radicals $-\text{O}-$ or $-\text{O}(\text{C}(\text{R}_6)_2)_s\text{O}-$;


R_6 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 1-6 carbon atoms, carboalkyl of 2-7 carbon atoms, carboxyalkyl (2-7 carbon atoms), phenyl, or phenyl optionally substituted with one or more halogen, alkoxy of 1-6 carbon atoms, trifluoromethyl, amino, alkylamino of 1-3 carbon atoms, dialkylamino of 2-6 carbon atoms, nitro, cyano, azido, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, carboxyl, carboalkoxy of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, or alkyl of 1-6 carbon atoms;

R_2 , is selected from the group consisting of


5

- 149 -


10




15




20




25



30



35




40

45


50

55


- 150 -



R₃ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,






with the proviso that at least one of the R₃ groups is selected from the group



- 151 -

R₅ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,



R₈, and R₉ are each, independently, $-(\text{C}(\text{R}_6)_2)_r\text{NR}_6\text{R}_6$, or $-(\text{C}(\text{R}_6)_2)_r\text{OR}_6$;

J is independently hydrogen, chlorine, fluorine, or bromine;

Q is alkyl of 1-6 carbon atoms or hydrogen;

a = 0 or 1;

g = 1-6;

k = 0-4;

n is 0-1;

p = 2-4;

q = 0-4;

r = 1-4;

s = 1-6;

u = 0-4 and v = 0-4, wherein the sum of u+v is 2-4;

or a pharmaceutically acceptable salt thereof,

provided that

when R₆ is alkenyl of 2-7 carbon atoms or alkynyl of 2-7 carbon atoms, such

alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

and further provided that

when Y is -NR₆- and R₇ is -NR₆R₆, -N(R₆)₃⁺, or -NR₆(OR₆), then g = 2-6;

- 152 -

when M is -O- and R₇ is -OR₆, then p = 1-4;


when Y is -NR₆-, then k = 2-4;

when Y is -O- and M or W is -O-, then k = 1-4

when W is not a bond with Het bonded through a nitrogen atom, then q = 2-4

5 and when W is a bond with Het bonded through a nitrogen atom and Y is -O- or -NR₆-, then k = 2-4.

9. A pharmaceutical composition which comprises a compound of formula 1 having the structure



wherein:

X is cycloalkyl of 3 to 7 carbon atoms, which may be optionally substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a pyridinyl, pyrimidinyl, or phenyl ring wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally mono-, di-, or tri-substituted with a substituent selected from the group consisting of halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, carboxyalkyl of 2-7 carbon atoms, carboalkoxyalkyl of 3-8 carbon


- 153 -

atoms, aminoalkyl of 1-5 carbon atoms, N-alkylaminoalkyl of 2-9 carbon atoms, N,N-dialkylaminoalkyl of 3-10 carbon atoms, N-alkylaminoalkoxy of 2-9 carbon atoms, N,N-dialkylaminoalkoxy of 3-10 carbon atoms, mercapto, and benzoylamino;

5 Z is -NH-, -O-, -S-, or -NR- ;

15 R is alkyl of 1-6 carbon atoms, or carboalkyl of 2-7 carbon atoms;

G₁, G₂, R₁, and R₄ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, alkenyloxy of 2-6 carbon atoms, alkynyloxy of 2-6 carbon atoms, hydroxymethyl, halomethyl, alkanoyloxy of 1-6 carbon atoms, alkenoyloxy of 3-8 carbon atoms, alkynoyloxy of 3-8 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkenoyloxymethyl of 4-9 carbon atoms, alkynoyloxymethyl of 4-9 carbon atoms, alkoxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, alkylsulphinyl of 1-6 carbon atoms, alkylsulphonyl of 1-6 carbon atoms, alkylsulfonamido of 1-6 carbon atoms, alkenylsulfonamido of 2-6 carbon atoms, alkynylsulfonamido of 2-6 carbon atoms, hydroxy, trifluoromethyl, trifluoromethoxy, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzyl, amino, hydroxyamino, alkoxyamino of 1-4 carbon atoms, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, N-alkylcarbamoyl, N,N-dialkylcarbamoyl, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, phenylamino, benzylamino,



25 $R_7-(C(R_6)_2)_q-Y- , R_7-(C(R_6)_2)_p-M-(C(R_6)_2)_k-Y- ,$ or $Het-(C(R_6)_2)_q-W-(C(R_6)_2)_k-Y-$

with the proviso that either G₁ or G₂ or both G₁ and G₂ must be a radical selected from the group

5

10

15

20

10

25

15

30

35

20

40




45

25


50

55

- 154 -



Y is a divalent radical selected from the group consisting of




R_7 is $-NR_6R_6$, $-J$, $-OR_6$, $-N(R_6)_3^+$, or $-NR_6(OR_6)$;

R'_7 is $-NR_6(OR_6)$, $-N(R_6)_3^+$, alkenoxy of 1-6 carbon atoms, alkynoxy of 1-6 carbon atoms, N-alkyl-N-alkenylamino of 4 to 12 carbon atoms, N,N-dialkenylamino of 6-12 carbon atoms, N-alkyl-N-alkynylamino of 4 to 12 carbon atoms, N-alkenyl-N-alkynylamino of 4 to 12 carbon atoms, or N,N-dialkynylamino of 6-12 carbon atoms with the proviso that the alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

M is $>NR_6$, $-O-$, $>N-(C(R_6)_2)_pNR_6R_6$, or $>N-(C(R_6)_2)_p-OR_6$;

W is $>NR_6$, $-O-$ or is a bond;

Het is a heterocycle selected from the group consisting of morpholine, thiomorpholine, thiomorpholine S-oxide, thiomorpholine S,S-dioxide, piperidine, pyrrolidine, aziridine, pyridine, imidazole, 1,2,3-triazole, 1,2,4-triazole, thiazole, thiazolidine, tetrazole, piperazine, furan, thiophene, tetrahydrothiophene, tetrahydrofuran, dioxane, 1,3-dioxolane,



- 155 -

wherein the heterocycle is optionally mono- or di-substituted on carbon or nitrogen with R_6 , optionally mono- or di-substituted on carbon with hydroxy, $-N(R_6)_2$, or $-OR_6$, optionally mono or di-substituted on carbon with the mono-valent radicals $-(C(R_6)_2)_sOR_6$ or $-(C(R_6)_2)_sN(R_6)_2$, or optionally mono or di-substituted on a saturated carbon with divalent radicals $-O-$ or $-O(C(R_6)_2)_sO-$;


R_6 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 1-6 carbon atoms, carboalkyl of 2-7 carbon atoms, carboxyalkyl (2-7 carbon atoms), phenyl, or phenyl optionally substituted with one or more halogen, alkoxy of 1-6 carbon atoms, trifluoromethyl, amino, alkylamino of 1-3 carbon atoms, dialkylamino of 2-6 carbon atoms, nitro, cyano, azido, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, carboxyl, carboalkoxy of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, or alkyl of 1-6 carbon atoms;

R_2 is selected from the group consisting of


5

- 156 -


10




15




20




25



30



35






40



45

50



55



30






10



- 158 -

R₅ is independently hydrogen, alkyl of 1-6 carbon atoms, carboxy, carboalkoxy of 1-6 carbon atoms, phenyl, carboalkyl of 2-7 carbon atoms,



R₈, and R₉ are each, independently, -(C(R₆)₂)_rNR₆R₆, or -(C(R₆)₂)_rOR₆;

J is independently hydrogen, chlorine, fluorine, or bromine;

Q is alkyl of 1-6 carbon atoms or hydrogen;

a = 0 or 1;

g = 1-6;

k = 0-4;

n is 0-1;

p = 2-4;

q = 0-4;

r = 1-4;

s = 1-6;

u = 0-4 and v = 0-4, wherein the sum of u+v is 2-4;

or a pharmaceutically acceptable salt thereof,

provided that

when R₆ is alkenyl of 2-7 carbon atoms or alkynyl of 2-7 carbon atoms, such alkenyl or alkynyl moiety is bound to a nitrogen or oxygen atom through a saturated carbon atom;

and further provided that

when Y is -NR₆- and R₇ is -NR₆R₆, -N(R₆)₃⁺, or -NR₆(OR₆), then g = 2-6;

when M is -O- and R₇ is -OR₆, then p = 1-4;

- 159 -

when Y is $-NR_6-$, then $k = 2-4$;


when Y is $-O-$ and M or W is $-O-$, then $k = 1-4$;

when W is not a bond with Het bonded through a nitrogen atom, then $q = 2-4$;

and when W is a bond with Het bonded through a nitrogen atom and Y is $-O-$ or $-NR_6-$, then $k = 2-4$.

10. A process for preparing a compound of formula 1 or a pharmaceutically acceptable salt thereof which comprises

(a) reacting a compound having the formula



where R₁, G₁, G₂, R₄, Z, n and X are as defined above with a dehydrating agent so as to convert the aminocarbonyl group into a cyano group, or

(b) reacting a compound having the formula



or a salt thereof with a compound having the formula



where Q is a leaving group and A₁, A₂ and A₃ are such that A₁-NA₂-A₃ is a compound conforming with formula 1; or

(c) reacting a compound having the formula



or a salt thereof with a compound having the formula



where Q is as defined above and A₄ and A₅ are such that A₄-O-A₅ is a compound conforming with formula 1; or

(d) adding an acid to a compound having formula 1 so that an acid addition salt is prepared.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/22056

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/54 A61K31/47 C07D401/12 C07D417/12 C07D215/56
C07D401/04 C07D491/10 C07D405/12 C07D413/12
/(C07D491/10,317:00,211:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 02437 A (GLAXO GROUP LTD.) 22 January 1998 (1998-01-22) page 1; claim 1 ---	1,9
A	EP 0 787 722 A (AMERICAN CYANAMID COMPANY) 6 August 1997 (1997-08-06) page 3, line 1 - line 8; claim 1 ---	1,9
A	WO 96 09294 A (THE WELLCOME FOUNDATION LTD.) 28 March 1996 (1996-03-28) page 5; claim 1 ---	1,9
P,X	WO 98 43960 A (AMERICAN CYANAMID COMPANY) 8 October 1998 (1998-10-08) complete document -----	1,9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 January 2000

Date of mailing of the international search report

26/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/22056

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 6-8
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 6-8
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/22056

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9802437 A	22-01-1998	AU 3693697 A	09-02-1998
		CA 2260058 A	22-01-1998
		CN 1230185 A	29-09-1999
		CZ 9900087 A	16-06-1999
		EP 0912572 A	06-05-1999
		NO 990123 A	13-01-1999
		PL 331154 A	21-06-1999
EP 787722 A	06-08-1997	AU 1252897 A	14-08-1997
		BR 9700850 A	01-09-1998
		CA 2196640 A	06-08-1997
		CZ 9700306 A	15-10-1997
		HU 9700344 A	28-10-1997
		JP 9221478 A	26-08-1997
		NO 970501 A	06-08-1997
		NZ 314184 A	28-10-1998
		SK 14497 A	10-09-1997
		US 5760041 A	02-06-1998
		ZA 9700913 A	04-08-1998
WO 9609294 A	28-03-1996	AU 3482495 A	09-04-1996
		EP 0782570 A	09-07-1997
		JP 10505600 T	02-06-1998
		TR 960233 A	21-06-1996
WO 9843960 A	08-10-1998	AU 6877798 A	22-10-1998
		NO 994798 A	24-11-1999